

CORRECTION

Correction: The effect of glutamine therapy on outcomes in critically ill patients: a meta-analysis of randomized controlled trials

Qi-Hong Chen¹, Yi Yang¹, Hong-Li He¹, Jian-Feng Xie¹, Shi-Xia Cai¹, Ai-Ran Liu¹, Hua-Ling Wang² and Hai-Bo Qiu^{1*} See related research by Chen *et al.*, http://ccforum.com/content/18/1/R8

After publication of our article in a recent issue of *Critical Care* [1], inconsistencies were identified in Figure 6. Four trials - by Fuentes-Orozco and colleagues [2,3], Hall and colleagues [4], and Goeters and colleagues [5] - were allocated to

the wrong subgroup in our meta-analysis. They should have been included in the first subgroup (glutamine <0.3 g/kg per day); this does not change the significance of the results (P = 0.01). The correct Figure 6 is given here in full as Figure 1.




Figure 1 A subgroup meta-analysis of the effects of different dosages of glutamine on mortality in critically ill patients (fixed effects models). CI, confidence interval; M-H, Mantel-Haenszel.

¹Department of Critical Care Medicine, Zhong-Da Hospital, School of Medicine, Southeast University, 87 Dingjiaqiao Road, Nanjing 210009, P.R. China Full list of author information is available at the end of the article



^{*} Correspondence: haiboq2000@gmail.com

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Critical Care Medicine, Zhong-Da Hospital, School of Medicine, Southeast University, 87 Dingjiaqiao Road, Nanjing 210009, P.R. China. ²Department of Critical Care Medicine, Su-Bei Hospital of Jiangsu Provience & Clinical Medical School, Yangzhou University, Yangzhou, Jiangsu, P.R. China.

Published: 17 June 2014

References

- Chen QH, Yang Y, He HL, Xie JF, Cai SX, Liu AR, Wang HL, Qiu HB: The effect of glutamine therapy on outcomes in critically ill patients: a meta-analysis of randomized controlled trials. Crit Care 2014, 18:R8.
- Fuentes-Orozco C, Anaya-Prado R, González-Ojeda A, Arenas-Márquez H, Cabrera-Pivaral C, Cervantes-Guevara G, Barrera-Zepeda LM: L-alanyl-Lglutamine-supplemented parenteral nutrition improves infectious morbidity in secondary peritonitis. Clin Nutr 2004, 23:13–21.
- Fuentes-Orozco C, Cervantes-Guevara G, Muciño-Hernández I, López-Ortega A, Ambriz-González G, Gutiérrez-de-la-Rosa JL, Gómez-Herrera E, Hermosillo-Sandoval JM, González-Ojeda A: L-alanyl-L-glutamine-supplemented parenteral nutrition decreases infectious morbidity rate in patients with severe acute pancreatitis. JPEN J Parenter Enteral Nutr 2008, 32:403-411.
- Hall JC, Dobb G, Hall J, de Sousa R, Brennan L, McCauley R: A prospective randomized trial of enteral glutamine in critical illness. *Intensive Care Med* 2003, 29:1710–1716.
- Goeters C, Wenn A, Mertes N, Wempe C, Van Aken H, Stehle P, Bone HG: Parenteral L-alanyl-L-glutamine improves 6-month outcome in critically ill patients. Crit Care Med 2002, 30:2032–2037.

doi:10.1186/cc13926

Cite this article as: Chen *et al.*: Correction: The effect of glutamine therapy on outcomes in critically ill patients: a meta-analysis of randomized controlled trials. *Critical Care* 2014 **18**:436.