ERRATUM Open Access



Erratum to: Cardiac dysfunction induced by weaning from mechanical ventilation: incidence, risk factors, and effects of fluid removal

Jinglun Liu^{1,2,3,4†}, Feng Shen^{1,2,3,5†}, Jean-Louis Teboul^{1,2,3}, Nadia Anguel^{1,2,3}, Alexandra Beurton^{1,2,3}, Nadia Bezaz^{1,2,3}, Christian Richard^{1,2,3} and Xavier Monnet^{1,2,3*}

Erratum

Unfortunately this article was published with an error. During production of the original article [1], "with WiPO" and "without WiPO" on " Δ CI% during PLR" has been switched on Fig. 3. The correct figure is shown below.

Author details

¹Université Paris-Sud, Faculté de Médecine, Université Paris-Saclay, Le Kremlin-Bicêtre, France. ²AP-HP, Service de réanimation médicale, Hôpital de Bicêtre, 78, rue du Général Leclerc, 94 270 Le Kremlin-Bicêtre, France. ³Inserm UMR_S 999, Hôpital Marie Lannelongue, Le Plessis-Robinson, France. ⁴Department of Emergency Medicine and Critical Care Medicine, The First Affiliated Hospital of Chongqing Medical University, Chongqing, China. ⁵Department of Critical Care Medicine, Affiliated Hospital of Guizhou Medical University, Guiyang, China.

Published online: 08 March 2017

Reference

 Monnet X. Cardiac dysfunction induced by weaning from mechanical ventilation: incidence, risk factors, and effects of fluid removal. BMC Critical Care. 2016;20:369. doi:10.1186/s13054-016-1533-9.

²AP-HP, Service de réanimation médicale, Hôpital de Bicêtre, 78, rue du Général Leclerc, 94 270 Le Kremlin-Bicêtre, France Full list of author information is available at the end of the article



^{*} Correspondence: xavier.monnet@aphp.fr

[†]Equal contributors

¹Université Paris-Sud, Faculté de Médecine, Université Paris-Saclay, Le Kremlin-Bicêtre, France




Fig. 3 Changes in haemodynamic variables, plasma protein concentration, and extravascular lung water during the spontaneous breathing trial (SBT). *p < 0.05 at the end of SBT vs. before SBT; *p < 0.05 cases without WiPO vs. cases with WiPO. CI cardiac index, EVLW extravascular lung water, PLR passive leg raising, SBP systolic arterial blood pressure, WiPO weaning-induced pulmonary oedema