

LETTER

ROTEM®-guided coagulation factor concentrate therapy in trauma: 2-year experience in Venice, Italy

Alberto Grassetto*, Marco De Nardin, Bernadetta Ganzerla, Monica Geremia, Debora Saggiaro, Elena Serafini, Silvia Zampieri, Manuela Toffoli, Daniele Penzo, Antonio Bossi and Carlo Maggiolo

Haemostatic therapy for trauma-induced coagulopathy is typically based on administration of allogeneic blood products, although the evidence supporting this approach is poor [1]. Fixed-ratio protocols have been proposed for administering fresh frozen plasma, red blood cells and platelets, but the optimal ratio has not been established and the speed of intervention may be more important [2].

In contrast to fixed-ratio treatment, coagulation factor concentrate therapy guided by point-of-care monitoring allows patients' actual needs to be targeted [3]. Our initial experience with ROTEM® (Tem International GmbH, Munich, Germany) indicated correlation between the clinical condition and extent of coagulopathy, suggesting a need for the early identification and treatment of coagulopathy that ROTEM® enables. Altogether, these factors provided a rationale for implementing ROTEM®-guided therapy for trauma patients in our hospital.

In our experience, this approach is feasible and can replace formula-driven treatment. We found that coagulation factor concentrates (fibrinogen concentrate and prothrombin complex concentrate) correct coagulopathy effectively and rapidly, indicated by normalisation of ROTEM® parameters among bleeding trauma patients (Table 1). European guidelines for managing trauma raised the target fibrinogen concentration to 1.5 to 2 g/l [4], which we find difficult to reach without using fibrinogen concentrate. Without a comparator group, our data are insufficient to show reduced red blood cell transfusion or improvements in morbidity/mortality. However, we did see a progressive reduction in fresh frozen plasma consumption. Another advantage of using a ROTEM®-guided approach is the opportunity to detect hyperfibrinolysis. As reported elsewhere, we found that fulminant hyperfibrinolysis is associated with high mortality. Fulminant hyperfibrinolysis may potentially be considered the last gasp of the coagulation system; it may be a marker not only of severe coagulopathy, but also of

poor clinical outcome. Our experience also suggests that patients with massive bleeding may benefit from immediate, proactive administration of 1 g tranexamic acid followed by 2 to 4 g fibrinogen concentrate, with further doses as soon as ROTEM® results are available.

Fibrinogen concentrate is currently imported in Italy and we use it according to the manufacturer's label. In some countries the product is licensed only for congenital deficiency. However, it is possible to use life-saving drugs for indications beyond the label, providing the physician is convinced that this use is in the patient's best interest; such practice is regulated by health authorities in several countries. High-quality, randomised controlled trials are lacking for both allogeneics and coagulation factor concentrates in trauma, creating a degree of uncertainty with both of these options. Nevertheless, we consider the rationale to be stronger for ROTEM®-guided, concentrate-based therapy.

Competing interests

The authors declare that they have no competing interests.

Acknowledgements

Editorial assistance was provided by medical writers from Meridian HealthComms during the preparation of this manuscript. Financial support for this assistance was provided by CSL Behring.

Published: 21 May 2012

References

1. Yang L, Stanworth S, Hopewell S, Doree C, Murphy M: Is fresh-frozen plasma clinically effective? An update of a systematic review of randomized controlled trials. *Transfusion* 2012. [Epub ahead of print]
2. Riskin DJ, Tsai TC, Riskin L, Hernandez-Boussard T, Purtil M, Maggio PM, Spain DA, Brundage SI: Massive transfusion protocols: the role of aggressive resuscitation versus product ratio in mortality reduction. *J Am Coll Surg* 2009, **209**:198-205.
3. Schöch H, Maegele M, Solomon C, Görlinger K, Voelckel W: Early and individualized goal-directed therapy for trauma-induced coagulopathy. *Scand J Trauma Resusc Emerg Med* 2012, **20**:15.
4. Rossaint R, Bouillon B, Cerny V, Coats TJ, Duranteau J, Fernandez-Mondejar E, Hunt BJ, Komadina R, Nardi G, Neugebauer E, Ozier Y, Riddez L, Schultz A, Stahel PF, Vincent JL, Spahn DR: Management of bleeding following major trauma: an updated European guideline. *Crit Care* 2010, **14**:R52.

*Correspondence: alberto.grassetto@gmail.com

Dipartimento Emergenza Urgenza, UOC Anestesia e Rianimazione, Ospedale dell'Angelo di Mestre, Via Paccagnella 11, 30174 Zelarino, Venice, Italy

Table 1. Details of bleeding trauma patients receiving ROTEM®-guided coagulation factor concentrate treatment

ISS on arrival	ROTEM® parameters on arrival				Haemostatic treatment administered in response to initial ROTEM® assessment					ROTEM® parameters following haemostatic treatment				Outcome
	EXTEM		FIBTEM		Prothrombin complex concentrate (IU)	Fibrinogen concentrate (g)	Tranexamic acid (g)	Platelet concentrate (U)	EXTEM		FIBTEM			
	CT (seconds)	A10 (mm)	MCF (mm)	MCF (mm)					CT (seconds)	A10 (mm)	MCF (mm)	MCF (mm)		
1	25	130	31	44	3	2,000	6	2	0	65	42	54	12	Survived
2	25	75	37	48	5	0	2	0	0	N/A	N/A	N/A	N/A	Survived
3	34	59	54	62	10	0	1	0	0	N/A	N/A	N/A	N/A	Survived
4	43	70	35	43	3	0	3	0	0	40	32	34	10	Survived ^a
5	34	95	40	51	7	500	2	0	0	62	42	54	8	Survived ^a
6	25	167	18	29	2	1,000	5	2	2	62	39	50	10	Survived
7	66 ^b	216 ^c	14 ^c	22 ^c	0	1,000	7	2	2	59	38	51	13	Died ^d
8	66 ^b	139 ^e	22 ^c	34 ^c	0	2,000	8	2	2	74	26	38	8	Died ^e
9	34	71	46	56	10	0	3	0	0	61	56	65	14	Survived
10	27	170	20	32	3	1,500	6	2	2	50	39	50	12	Survived
11	45	102	28	37	5	1,500	3	2	1	48	53	63	12	Died ^f
12	36	114	20	29	5	1,500	4	2	1	61	39	48	9	Survived ^a

A10, clot amplitude (firmness) 10 minutes after clotting time (CT); EXTEM, extrinsically activated thromboelastometric test; FIBTEM, thromboelastometric test of fibrin-based clotting; ISS, injury severity score; IU, international units; MCF, maximum clot firmness; N/A, not available. ^aPatient received platelets following the second ROTEM® assessment. ^bFulminant hyperfibrinolysis was detected, so aprotinin-controlled thromboelastometric test (APTEM) parameters were initially used to guide treatment. Both patients also received fresh frozen plasma after the second ROTEM® assessment. ^cValue obtained using the APTEM assay. ^dCause of death was multiple organ failure. ^eCause of death was abdominal aortic rupture and unstopable haemorrhage. ^fCause of death was brain death.