

COMMENTARY

Early vs late start of dialysis: it's all about timing

Etienne Macedo and Ravindra L Mehta*

See related research by Shiao *et al.*, <http://ccforum.com/content/13/5/R171>

Abstract

Acute kidney injury (AKI) is now well recognized as an independent risk factor for increased morbidity and mortality, particularly when dialysis is needed. The wide variation in dialysis utilization contributes to a lack of consensus on what parameters should guide the decision to start dialysis. While the association of early initiation of dialysis with survival benefit was first demonstrated four decades ago, few studies in the modern era of dialysis have addressed time of dialysis initiation. Though listed as one of the top priorities in research on AKI, timing of dialysis initiation has not been included as a factor in any of the large, randomized controlled trials in this area.

Whether or not to provide dialytic support and when to start are two dilemmas for clinicians managing patients with a sudden decline in renal function. Earlier initiation is thought to be associated with better control of uremia, acidemia, electrolyte imbalances, and volume accumulation. However, the appreciation of the effect of time of initiation depends on what is considered early versus late. Various studies have considered early versus late time of dialysis initiation based on arbitrary thresholds of traditional serum biomarkers or time from intensive care unit (ICU) admission or from the diagnosis of acute kidney injury (AKI). The study by Shiao and colleagues [1] in a recent issue of *Critical Care* provides support that early start may be beneficial and offers an additional approach to identifying a starting point for dialysis.

Although a recent meta-analysis that included four randomized controlled trials and 19 observational studies conducted over four decades suggested that early dialysis initiation may have a beneficial effect on survival [2],

what constitutes early versus late has yet to be defined. Two main approaches have been used for stratifying early and late. In most studies, levels of solutes (blood urea nitrogen [BUN] and serum creatinine) have been used to define cutoffs for early and late dialysis initiation, showing variable results on different patient populations. In post-traumatic patients, BUN levels of less than 60 mg/dL at dialysis initiation were associated with a 20% absolute reduction in mortality [3]. Wu and colleagues [4] found a BUN level of less than 80 mg/dL to be predictive of mortality in patients requiring dialysis for acute liver failure after surgery. In the general ICU population, a large observational study (Program to Improve Care in Acute Renal Disease, or PICARD) showed an increased risk of mortality in patients with higher BUN concentrations (>76 mg/dL) [5]. However, a recent randomized single-center clinical trial in 106 critically ill patients with oliguric AKI [6] demonstrated that despite early dialysis at a BUN level of less than 48 mg/dL in comparison with 105 mg/dL for late dialysis, there was no difference in outcomes. These findings suggest that BUN levels are relatively insensitive as a target criterion for starting dialysis.

A second approach was used in the beginning and ending supportive therapy for the kidney (B.E.S.T. kidney) study, in which investigators included in the analysis a stratification of early or late based on time to initiate dialysis from ICU admission, besides the absolute urea and creatinine, and relative change in urea and creatinine [7]. Although absolute or delta BUN levels were insensitive in predicting mortality, the analysis by time from ICU admission showed a more than twofold increase in the odds of hospital mortality. However, in two recent, large, randomized controlled trials of dialysis dose, time to initiate dialysis was assessed from ICU admission and was not associated with outcomes [8,9].

Though using heterogeneous definitions of early initiation, these large observational cohorts and small randomized trials suggest that there may be a survival advantage to an early start for dialysis. They also highlight the need for better parameters to define the need for dialysis and the delineation of what is early and late. In the postoperative setting, the timing and type of renal

*Correspondence: rmehta@ucsd.edu
Department of Medicine, University of California at San Diego, 200 West Arbor Drive, MC 8342, San Diego, CA 92103, USA

insult are more homogenous, providing an opportunity to ascertain the benefits of earlier dialysis initiation when the event associated with AKI is known. Two cardiac surgery studies demonstrated a benefit in earlier initiation [10,11]. In these studies, urine output of less than 100 mL during the first 8 hours after bypass surgery was a criterion to initiate dialysis regardless of solute clearance. Mortality rates appeared to be dramatically reduced in both studies in the early dialysis groups. Similar findings were seen in a small study of 21 patients treated with prophylactic perioperative hemodialysis [12].

In the study of Shiao and colleagues [1], 98 patients who required dialysis in the postoperative period of abdominal surgery were categorized as early or late dialysis initiation based on the estimated glomerular filtration rate criteria of the RIFLE (Risk, Injury, Failure, Loss, and End-stage kidney disease) classification (simplified RIFLE, or sRIFLE). The earlier initiation group had lower ICU and hospital mortality rates than the late initiation group. These results suggest that the severity of renal injury may provide a better parameter than arbitrary values of traditional serum biomarkers (BUN and serum creatinine) for initiating dialysis. However, several questions still need to be answered. The RIFLE and Acute Kidney Injury Network classification systems are validated criteria for the severity of AKI but may not be the ideal parameters of early or late, as previously pointed out by Bellomo and colleagues [13]. The relationship of RIFLE classes at initiation and outcomes is subject to other influences that need to be considered. For instance, in the cohort of Shiao and colleagues, cardiac failure was an independent risk factor for in-hospital mortality. By their definition of cardiac failure (low cardiac output with a central venous pressure of greater than 12 mm Hg and a dopamine equivalent of greater than 5 µg/kg per minute), it is reasonable to assume that cardiac failure was a surrogate marker of fluid overload. This finding corroborates studies finding an inverse relationship between fluid accumulation and survival [14,15]. Additionally, other factors influence recognition of the severity of AKI. Shiao and colleagues found a lower prevalence of chronic kidney disease (CKD) in the late dialysis group, confirming data showing that an earlier identification of AKI among patients with prior CKD could modify the process of care delivered to these patients [16]. Thus, the time to recognize AKI, the severity and response to injury, and the contribution of non-renal factors may all influence the timing of initiation.

Timing of dialysis initiation is a potentially modifiable factor that may play an important role in determining patient outcomes. Based on current knowledge, we would recommend assessing patients for changes in renal function and using dialysis to support organ function and prevent complications rather than waiting for complete

renal shutdown prior to renal replacement [17]. Future research in this field is desperately needed and should include a combination of clinical and emerging biomarkers to inform these decisions. We look forward to doing away with comparisons of early versus late dialysis and focusing on improving outcomes with timely interventions of renal support individualized to patient need.

Abbreviations

AKI = acute kidney injury; BUN = blood urea nitrogen; CKD = chronic kidney disease; ICU = intensive care unit; RIFLE = Risk, Injury, Failure, Loss, and End-stage kidney disease.

Competing interests

The authors declare that they have no competing interests.

Published: 8 February 2010

References

1. Shiao CC, Wu VC, Li WY, Lin YF, Hu FC, Young GH, Kuo CC, Kao TW, Huang DM, Chen YM, Tsai PR, Lin SL, Chou NK, Lin TH, Yeh YC, Wang CH, Chou A, Ko WJ, Wu KD, the National Taiwan University Surgical Intensive Care Unit-Associated Renal Failure (NSARF) Study Group: **Late initiation of renal replacement therapy is associated with worse outcomes in acute kidney injury after major abdominal surgery.** *Crit Care* 2009, **13**:R171.
2. Seabra VF, Balk EM, Liangos O, Sosa MA, Cendoroglo M, Jaber BL: **Timing of renal replacement therapy initiation in acute renal failure: a meta-analysis.** *Am J Kidney Dis* 2008, **52**:272-284.
3. Gettings LG, Reynolds HN, Scalea T: **Outcome in post-traumatic acute renal failure when continuous renal replacement therapy is applied early vs. late.** *Intensive Care Med* 1999, **25**:805-813.
4. Wu VC, Ko WJ, Chang HW, Chen YS, Chen YW, Chen YM, Hu FC, Lin YH, Tsai PR, Wu KD: **Early renal replacement therapy in patients with postoperative acute liver failure associated with acute renal failure: effect on postoperative outcomes.** *J Am Coll Surg* 2007, **205**:266-276.
5. Liu KD, Himmelfarb J, Paganini E, Ikizler TA, Soroko SH, Mehta RL, Chertow GM: **Timing of initiation of dialysis in critically ill patients with acute kidney injury.** *Clin J Am Soc Nephrol* 2006, **1**:915-919.
6. Bouman CS, Oudemans-Van Straaten HM, Tijssen JG, Zandstra DF, Kesecioglu J: **Effects of early high-volume continuous venovenous hemofiltration on survival and recovery of renal function in intensive care patients with acute renal failure: a prospective, randomized trial.** *Crit Care Med* 2002, **30**:2205-2211.
7. Bagshaw SM, Uchino S, Bellomo R, Morimatsu H, Morgera S, Schetz M, Tan I, Bouman C, Macedo E, Gibney N, Tolwani A, Oudemans-van Straaten HM, Ronco C, Kellum JA; Beginning and Ending Supportive Therapy for the Kidney (BEST Kidney) Investigators: **Timing of renal replacement therapy and clinical outcomes in critically ill patients with severe acute kidney injury.** *J Crit Care* 2009, **24**:129-140.
8. VA/NIH Acute Renal Failure Trial Network, Palevsky PM, Zhang JH, O'Connor TZ, Chertow GM, Crowley ST, Choudhury D, Finkel K, Kellum JA, Paganini E, Schein RM, Smith MW, Swanson KM, Thompson BT, Vijayan A, Watnick S, Star RA, Peduzzi P: **Intensity of renal support in critically ill patients with acute kidney injury.** *N Engl J Med* 2008, **359**:7-20.
9. RENAL Replacement Therapy Study Investigators, Bellomo R, Cass A, Cole L, Finfer S, Gallagher M, Lo S, McArthur C, McGuinness S, Myburgh J, Norton R, Scheinkestel C, Su S: **Intensity of continuous renal-replacement therapy in critically ill patients.** *N Engl J Med* 2009, **361**:1627-1638.
10. Demirkılıç U, Kuralay E, Yenicesu M, Çağlar K, Oz BS, Cingöz F, Günay C, Yildirim V, Ceylan S, Arslan M, Vural A, Tatar H: **Timing of replacement therapy for acute renal failure after cardiac surgery.** *J Card Surg* 2004, **19**:17-20.
11. Elahi MM, Lim MY, Joseph RN, Dhannapunen RR, Spyt TJ: **Early hemofiltration improves survival in post-cardiotomy patients with acute renal failure.** *Eur J Cardiothorac Surg* 2004, **26**:1027-1031.
12. Durmaz I, Yagdi T, Calkavur T, Mahmudov R, Apaydin AZ, Posacioglu H, Atay Y, Engin C: **Prophylactic dialysis in patients with renal dysfunction undergoing on-pump coronary artery bypass surgery.** *Ann Thorac Surg* 2003, **75**:859-864.

13. Bellomo R, Kellum JA, Ronco C: **Comment on "RIFLE classification in patients with acute kidney injury in need of renal replacement therapy" by Maccariello et al.** *Intensive Care Med* 2007, **33**:1850; author reply 1851-1852.
14. Payen D, de Pont AC, Sakr Y, Spies C, Reinhart K, Vincent JL: **A positive fluid balance is associated with a worse outcome in patients with acute renal failure.** *Crit Care* 2008, **12**:R74.
15. Bouchard J, Soroko S, Chertow G, Himmelfarb J, Ikizler T, Paganini E, Mehta R: **Fluid accumulation, survival and recovery of kidney function in critically ill patients with acute kidney injury.** *Kidney Int* 2009, **76**:422-427.
16. Khosla N, Soroko SB, Chertow GM, Himmelfarb J, Ikizler TA, Paganini E, Mehta

- RL; Program to Improve Care in Acute Renal Disease (PICARD): **Preexisting chronic kidney disease: a potential for improved outcomes from acute kidney injury.** *Clin J Am Soc Nephrol* 2009, **4**:1914-1919.
17. Mehta RL: **Indications for dialysis in the ICU: renal replacement vs. renal support.** *Blood Purif* 2001, **19**:227-232.

doi:10.1186/cc8199

Cite this article as: Macedo E, Mehta RL: **Early vs late start of dialysis: it's all about timing.** *Critical Care* 2010, **14**:112.