

LETTER


Another explanation for decreased oxygen consumption in lactic acidosis

Viktor Rosival*

See related research by Protti *et al.*, <http://ccforum.com/content/14/1/R22>

In their recent paper, Protti and colleagues reported depressed oxygen consumption in patients with lactic acidosis due to biguanide intoxication and they suppose that the cause is inhibited mitochondrial respiration [1].

Another explanation for depressed oxygen consumption in these patients is also possible, however. If the blood pH is very low, glucose utilization is decreased [2] because the glycolytic enzyme phosphofructokinase is pH dependent – with decreasing pH, its activity is also decreasing [3]. Glucose utilization is an oxygen-consuming process:



The consequence of decreased utilization of glucose is thus also decreased oxygen consumption.

The patients reported by Protti and colleagues had on admission very low blood pH of 6.93 ± 0.20 and systemic oxygen consumption of 67 ± 28 ml/min/m² [1]. Systemic oxygen consumption ‘normalized within the next 48–72 hours’ and ‘Systemic O₂ consumption was positively associated with arterial pH’ ($P < 0.001$). According to Tables 2 and 3 [1], arterial pH reached normal values on days 2 to 3.

Depressed oxygen consumption in patients reported by Protti and colleagues can thus be explained by their very low blood pH.

Authors' response

Alessandro Protti and Luciano Gattinoni

We thank Dr Rosival for his stimulating comment.

Whether acidosis has an impact on oxygen consumption (VO₂) remains unclear. *In vitro*, several studies have demonstrated that tissue VO₂ only starts to diminish when the pH falls below 6 to 6.5 [4,5]. *In vivo*, both animal and clinical studies have reported normal, or even

increased, whole-body VO₂ during severe acidosis [6,7]. Accordingly, we have observed no correlation between VO₂ and arterial pH among 762 critically ill patients, at the time of admission to intensive care ($R^2 = 0.00$, $P = 0.88$ on linear regression analysis) [8].

In order to directly address the issue raised by Dr Rosival, we equipped two healthy, sedated and mechanically




Figure 1. Effect of metformin and lactic acid on arterial pH and oxygen consumption. Upper panel: data recorded from a pig infused with 8 g metformin (final serum drug concentration, 98 µg/ml). Lower panel: data recorded from a pig infused with lactic acid. Lactatemia equally increased from 1 to 25 mmol/l in the two animals. VO₂, oxygen consumption.

*Correspondence: rosival@hotmail.com
SYNLAB Department of Laboratory Medicine, Dérer's Hospital, Limbová 5, SK-833 05 Bratislava, Slovakia

ventilated pigs with a metabolic module (to record VO_2) and a pulmonary artery catheter (to compute the global oxygen delivery). Following baseline recordings, one animal received a continuous intravenous infusion of metformin whereas the other received lactic acid. Arterial pH, VO_2 and oxygen delivery were recorded hourly for 10 hours. As shown in Figure 1, metformin progressively decreased VO_2 but lactic acid did not. Changes in oxygen delivery were always minor.

We are thus tempted to believe that drug toxicity, rather than acidosis, was the major factor responsible for the decrease in VO_2 we observed in patients with biguanide-induced lactic acidosis.

Abbreviations

VO_2 , oxygen consumption.

Competing interests

The authors declare that they have no competing interests.

Published: 6 July 2010

References

1. Protti A, Russo R, Tagliabue P, Vecchio S, Singer M, Rudiger A, Foti G, Rossi A, Mistraretti G, Gattinoni L: **Oxygen consumption is depressed in patients with lactic acidosis due to biguanide intoxication.** *Crit Care* 2010, **14**:R22.
2. Van Nimmen D, Weyne J, Demeester G, Leusen I: **Local cerebral glucose utilization in systemic acidosis.** *Am J Physiol* 1984, **247**:R639-R645.
3. Trivedi B, Danforth WH: **Effect of pH on the kinetics of frog muscle phosphofructokinase.** *J Biol Chem* 1966, **241**:4110-4112.
4. Koehler AE, Reitzel RJ: **The effect of pH on the oxygen consumption of tissues.** *J Biol Chem* 1925, **64**:739-751.
5. Canzanelli A, Greenblatt M, Rogers GA, Rapport D: **The effect of pH on the in-vitro O_2 consumption of tissues.** *Am J Physiol* 1939, **127**:290-295.
6. Nahas GG, Ligou JC, Mehlman B: **Effects of pH changes on O_2 uptake and plasma catecholamine levels in the dog.** *Am J Physiol* 1960, **198**:60-66.
7. Fisher P, Kleinerman JI: **Total oxygen consumption and metabolic rate of patients with diabetic acidosis.** *J Clin Invest* 1952, **31**:126-130.
8. Gattinoni L, Brazzi L, Pelosi P, Latini R, Tognoni G, Pesenti A, Fumagalli R: **A trial of goal-oriented hemodynamic therapy in critically ill patients.** Sv O_2 Collaborative Group. *N Engl J Med* 1995, **333**:1025-1032.

doi:10.1186/cc9072

Cite this article as: Rosival V: **Another explanation for decreased oxygen consumption in lactic acidosis.** *Critical Care* 2010, **14**:427.