

CORRECTION

Open Access



Correction: The effect of treatment and clinical course during Emergency Department stay on severity scoring and predicted mortality risk in Intensive Care patients

Bart G. J. Candel^{1,2*}, Wouter Raven¹, Heleen Lameijer³, Wendy A. M. H. Thijssen⁴, Fabian Termorshuizen^{5,6,7}, Christiaan Boerma⁸, Nicolette F. de Keizer^{5,6,7}, Evert de Jonge⁹ and Bas de Groot¹

Correction to: Crit Care 26, 112 (2022).

<https://doi.org/10.1186/s13054-022-03986-2>

Following publication of the original article [1], the authors identified an error in the author name Fabian Termorshuizen. The family name contained an error.

The incorrect author name is: Fabian Temorshuizen.

The correct author name is: Fabian Termorshuizen.

The author group has been updated above and the original article [1] has been corrected.

Author details

¹Department of Emergency Medicine, Leiden University Medical Centre, Albinusdreef 2, 2300 RC Leiden, The Netherlands. ²Department of Emergency Medicine, Máxima Medical Centre, De Run 4600, 5504 DB Veldhoven, The Netherlands. ³Department of Emergency Medicine, Medical Centre Leeuwarden, Henri Dunantweg 2, 8934 AD Leeuwarden, The Netherlands. ⁴Department of Emergency Medicine, Catharina Hospital Eindhoven, Michelangelolaan 2, 5623 EJ Eindhoven, The Netherlands. ⁵Department of Medical Informatics, Amsterdam University Medical Center, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands. ⁶Amsterdam Public Health, Quality of Care,

Amsterdam, The Netherlands. ⁷National Intensive Care Evaluation (NICE) Foundation, Amsterdam, The Netherlands. ⁸Department of Intensive Care, Medical Center Leeuwarden, Henri Dunantweg 2, 8934 AD Leeuwarden, The Netherlands. ⁹Department of Intensive Care Medicine, Leiden University Medical Centre, Albinusdreef 2, 2300 RC Leiden, The Netherlands.

Published online: 11 May 2022

Reference

1. Candel BGJ, Raven W, Lameijer H, et al. The effect of treatment and clinical course during Emergency Department stay on severity scoring and predicted mortality risk in Intensive Care patients. *Crit Care*. 2022;26:112. <https://doi.org/10.1186/s13054-022-03986-2>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1186/s13054-022-03986-2>.

*Correspondence: bart.candel@mmc.nl

¹ Department of Emergency Medicine, Leiden University Medical Centre, Albinusdreef 2, 2300 RC Leiden, The Netherlands
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.