

CORRESPONDENCE

Open Access



Mechanisms of lung and diaphragmatic protection by high PEEP in obese COVID-19 ARDS: role of the body mass index

Joaquín Pérez^{1,2}, Javier Hernan Dorado¹, Emiliano Navarro^{2,3}, Caio C. A. Morais^{4,5} and Matías Accoce^{1,6,7*}

To the Editor:

Obesity increases the risk of requiring mechanical ventilation (MV) and developing acute respiratory distress syndrome (ARDS) in patients with coronavirus 2019 (COVID-19) [1].

Spontaneous breathing (SB) may potentially compromise lung and diaphragmatic protection in COVID-19-induced ARDS [2]. In recruitable models, high positive end-expiratory pressure (PEEP) was shown to render SB less injurious [3]. Nonetheless, the effects of high PEEP have been scarcely studied in clinical scenarios. Previous literature suggests that obese subjects might benefit from higher PEEP and these positive effects might be proportional to the obesity grade [4, 5].

We assessed whether the beneficial effects of high PEEP on lung and diaphragmatic protection depend on body mass index (BMI) in intubated obese patients recovering from COVID-19-associated ARDS.

This is an exploratory analysis of a physiological study conducted at Sanatorio Anchorena San Martín, Argentina. Consecutive adults with COVID-19-associated ARDS, BMI > 30 kg/m², intubation > 48 hs and SB in pressure support for > 2 hs were included.

The pressure support level was titrated by the attending physician. Esophageal pressure (P_{es}) was measured with an air-filled (1 mL) esophageal balloon (MBMed[®]-BA-A-008, non-latex) according to manufacturer

recommendations after balloon's correct positioning. Airway pressure, flow, volume and P_{es} were recorded by a dedicated system (FluxMed, MBMed[®]).

Our primary endpoints were the changes in muscular pressure (P_{mus}), pressure–time product per-minute (PTP_{min}) and dynamic driving transpulmonary pressure (ΔPL_{dyn}) from low-to-high PEEP and its correlation with BMI. Data were collected 10 min after changing PEEP (5–15cmH₂O). Respiratory variables averages were computed based on respiratory cycles of the last 30–60 s of each PEEP step considered stable. In 16/21 patients, chest wall elastance (E_{cw}) was assessed before SB was present. In the remaining, E_{cw} was measured applying a hyperventilation maneuver by increasing the pressure support until observing positive P_{es} deflections. All traces were analyzed using Biopac Student Lab PRO[®].

Correlations were performed with Spearman's rank test. To assess the differential effects of high PEEP, we analyzed subgroups based on a commonly used BMI cutoff (≥ 35 kg/m²) [4]. A linear mixed-effects regression model for repeated measures was fitted to assess interaction between PEEP (5–15cmH₂O) and BMI (≥ or < 35 kg/m²). *P* values were adjusted by post hoc Tukey's correction. A two-tailed *P* ≤ 0.05 was considered statistically significant. Data were analyzed with R4.0.3 software.

We studied 21 patients (age, median [IQR, 25th–75th]: 56 years [IQR, 51–67]; BMI: 34.3 kg/m² [IQR, 31.6–40.4]). At admission, 85% had moderate ARDS and 3 had mild ARDS. All the patients required neuromuscular blockers and 62% received prone ventilation.

At inclusion, the patients had 5 MV days [IQR, 3–7] and remained ventilated for additional 11 days [IQR, 5–20].

*Correspondence: matiasaccoce@gmail.com

¹ Intensive Care Unit, Sanatorio Anchorena San Martín, Provincia de Buenos Aires, Perdriel 4189, Villa Lynch, San Martín, Argentina
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

The $\text{PaO}_2/\text{FiO}_2$ was 214 mmHg [IQR, 183–252], pressure support was $8\text{cmH}_2\text{O}$ [IQR, 5–10], and inspired oxygen was 0.40 [IQR, 0.35–0.50]; the Richmond Agitation–Sedation Scale was –3 to –4.

The BMI significantly correlated with changes in P_{mus} , PTP_{min} and $\Delta\text{PL}_{\text{dyn}}$ (Fig. 1-A). We found an interaction between BMI and PEEP regarding to changes in P_{mus} ($F_{(1,19)}=6.2; P=0.022$), PTP_{min} ($F_{(1,19)}=7.4, P=0.013$) and $\Delta\text{PL}_{\text{dyn}}$ ($F_{(1,19)}=10.4, P=0.004$). Only the subgroup with $\text{BMI} \geq 35$ reduced the P_{mus} (mean difference (95%CI): $-3.3\text{cmH}_2\text{O}$ (-1.5 to -6.1)), PTP_{min} ($-94.5\text{cmH}_2\text{O.s/min}$ (-23.3 to -165.7)) and $\Delta\text{PL}_{\text{dyn}}$ ($-3.2\text{cmH}_2\text{O}$ (-1.6 to -5.7)) with high PEEP (Fig. 1-B). The results remained significant after excluding an outlier from the severely obese subgroup. In these patients ($\text{BMI} \geq 35$), the end-expiratory transpulmonary pressure (PL_{exp}) was more negative at low PEEP and became closer to $0\text{cmH}_2\text{O}$ at high PEEP (Fig. 1-C). The optimization of lung–diaphragmatic protection was explained by reduction of PTP-associated components, improvement in dynamic lung compliance (CL_{dyn}), neuromuscular ventilatory coupling and respiratory drive (Fig. 1-D).

We demonstrate that the obesity grade greatly influences the beneficial effects of PEEP on relevant monitoring variables directly linked to the lung and diaphragmatic protection.

In obese subjects, the increased pleural pressure may reduce end-expiratory lung volume and become PL_{exp} negative, causing atelectasis and airway closure [4, 5].

Alveolar and airways collapse may alter ventilation distribution and cause occult pendelluft, resulting in local/global overdistension [3]. This might impose higher loads on the respiratory muscles, jeopardizing the SB even more [5].

Our findings indicate that these pathological mechanisms may play a role during SB in obese COVID-19 ARDS patients and can be counteracted by PEEP [4, 5]. Nonetheless, despite fulfilling obesity criteria ($\text{BMI} > 30\text{kg/m}^2$), not all patients benefited from high PEEP. Accordingly, the least obese subgroup did not experience work of breathing unloading and deteriorated lung mechanics, neuroventilatory coupling and respiratory drive, suggesting that lung overdistension was probably induced. In this context, we suggest that P_{es} should be used to titrate PEEP during assisted ventilation to avoid the potential harms associated with unnecessarily high PEEP levels, taking a slightly positive PL_{exp} as a reasonable target [4].

Our study is mainly limited by the small sample size and the lack of direct measurements of lung–diaphragmatic injury. Additionally, PEEP steps were not random and the time spent during each PEEP level was short.

In summary, considering the complexity of SB in COVID-19-induced ARDS [2], high PEEP might be considered as a potentially useful tool to provide lung and diaphragmatic protection particularly in selected individuals with $\text{BMI} \geq 35\text{kg/m}^2$, where the obesity-associated mechanical alterations might be specially exacerbated.

(See figure on next page.)

Fig. 1 **A** Correlation between BMI and changes (%) in P_{mus} , PTP_{min} , and $\Delta\text{PL}_{\text{dyn}}$; **B** P_{mus} (upper), PTP_{min} (middle) and $\Delta\text{PL}_{\text{dyn}}$ (lower) with PEEP 5 and $15\text{cmH}_2\text{O}$; **C** end-expiratory transpulmonary pressure at PEEP 5 and $15\text{cmH}_2\text{O}$; **D** mean (95% CI) percentage of change in PTP-associated components (intrinsic PEEP, resistance, elastance), dynamic lung compliance (CL_{dyn}), neuromuscular ventilatory coupling expressed as tidal volume-to-pressure–time product per breath ($\text{VT}/\text{PTP}_{\text{breath}}$) and intensity of respiratory drive expressed as esophageal pressure to neural inspiratory time (P_{es}/Ti). In panels **B**, **C** and **D**, the patients are divided in subgroups according to $\text{BMI} < 35\text{kg/m}^2$ ($n = 12$; median BMI 31.8 [IQR, 30.7 – 33.3] kg/m^2) and $\geq 35\text{kg/m}^2$ ($n = 9$, median BMI 43.6 [IQR, 38.5 – 48.9] kg/m^2)




Fig. 1 (See legend on previous page.)

Abbreviations

MV: Mechanical ventilation; ARDS: Acute respiratory distress syndrome; COVID-19: Coronavirus disease 2019; SB: Spontaneous breathing; PEEP: Positive end-expiratory pressure; BMI: Body mass index; P_{es} : Esophageal pressure; P_{mus} : Muscular pressure; PTP_{min} : Pressure–time product per minute; ΔP_{L-dyn} : Peak dynamic driving transpulmonary pressure; PaO_2/FiO_2 : Arterial oxygen pressure to inspired oxygen; P_{es}/Ti : Esophageal pressure to neural inspiratory time; VT/PTP_{breath} : Tidal volume-to-pressure–time product per breath; CL_{dyn} : Dynamic lung compliance; n.s: No statistically significant; IQR: Interquartile range.

Acknowledgements

The authors would like to thank Lic. Daniela Ines Gilgado, Lic. Gimena Paola Cardoso and all the staff of the Intensive Care Unit of Sanatorio Anchorena San Martín that made it possible to carry out this investigation.

Author contributions

J.P., J.H.D., M.A. and E.N. contributed to the study concept and design. J.P., J.H.D. and M.A. contributed to the acquisition of data. J.P., J.H.D., M.A., E.N. and C.C.A.M. contributed to the analysis and interpretation of data. J.P., J.H.D., M.A., E.N. and C.C.A.M. contributed to drafting the manuscript and critically revising the manuscript for important intellectual content. All authors read and approved the final manuscript.

Funding

There was no financial funding.

Availability of data and materials

The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

The original protocol was approved by the Institutional review board (approval #08-2019) and registered in clinicaltrials.gov (NCT04524091). The informed consent was obtained from the patient's next of kin.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Intensive Care Unit, Sanatorio Anchorena San Martín, Provincia de Buenos Aires, Perdriel 4189, Villa Lynch, San Martín, Argentina. ²Intensive Care Unit, Hospital de Agudos Carlos G. Durand, Ciudad Autónoma de Buenos Aires, Argentina. ³Centro del Parque, Ciudad Autónoma de Buenos Aires, Argentina. ⁴Divisão de Pneumologia Instituto Do Coração (Incor), Hospital das Clínicas Faculdade de Medicina, Universidade de São Paulo, São Paulo, Brazil. ⁵Departamento de Fisioterapia, Universidade Federal de Pernambuco, Recife, Brazil. ⁶Intensive Care Unit, Hospital de Quemados "Arturo Umberto Illia", Ciudad Autónoma de Buenos Aires, Argentina. ⁷Universidad Abierta Interamericana. Facultad de Medicina Y Ciencias de La Salud, Buenos Aires, Argentina.

Received: 9 March 2022 Accepted: 7 June 2022

Published online: 17 June 2022

References

- Kompaniyets L, Goodman AB, Belay B, et al. Body mass index and risk for COVID-19–related hospitalization, intensive care unit admission, invasive mechanical ventilation, and death—United States, March–December 2020. *MMWR Morb Mortal Wkly Rep.* 2021;70:355–61.
- Esnault P, Cardinale M, Hraiech S, Goutorbe P, Baumstrack K, Prud'homme E, et al. High respiratory drive and excessive respiratory efforts predict relapse of respiratory failure in critically ill patients with COVID-19. *Am J Respir Crit Care Med.* 2020;202(8):1173–8.
- Morais CCA, Koyama Y, Yoshida T, Plens GM, Gomes S, Lima CAS, et al. High positive end-expiratory pressure renders spontaneous effort non-injurious. *Am J Respir Crit Care Med.* 2018;197(10):1285–96.
- De Santis Santiago R, Teggie Droghi M, Fumagalli J, Marrazzo F, Florio G, Grassi LG, et al. Lung rescue team investigators. High pleural pressure prevents alveolar overdistension and hemodynamic collapse in ARDS with class III obesity. *Am J Respir Crit Care Med.* 2020;203(5):575–84.
- Teggie Droghi M, De Santis Santiago RR, Pinciroli R, Marrazzo F, Bittner EA, Amato MBP, Kacmarek RM, Berra L. High positive end-expiratory pressure allows extubation of an obese patient. *Am J Respir Crit Care Med.* 2018;198(4):524–5.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

