

CORRESPONDENCE

Open Access



Optimal PEEP with lowest (least injurious) transpulmonary driving pressure can be determined by a rapid two-PEEP-step procedure without esophageal pressure

O. Stenqvist*

To the Editor,

In the recent study on personalized optimal PEEP in hypoxemic patients during pressure support ventilation (PSV), Slobod and coworkers assessed collapse and overdistension by electric impedance tomography (EIT) during a PEEP trial [1]. Normally, this assessment is based on airway driving pressure (ΔPAW) and respiratory system compliance, but in this study, lung compliance (CL) and transpulmonary pressure by esophageal pressure were the basis of the analysis. They showed that the PEEP level with a balance between collapse and overdistension tended to coincide with the level where *lung* compliance is highest. Consequently, transpulmonary driving pressure (ΔPL) was lowest at this PEEP level, requiring the lowest inspiratory effort during PSV.

The authors should be commended for the important step to exchange respiratory system compliance and airway driving pressure for *lung* compliance and *transpulmonary* driving pressure because transpulmonary pressure is the pressure that directly affects lung tissue. In addition, it has been shown that the PEEP level with lowest airway driving pressure not necessarily coincides

with the PEEP level with lowest transpulmonary driving pressure [2, 3].

Protective ventilation requires assessment of lung mechanics and optimal PEEP as early as possible after start of mechanical ventilation. It is unlikely that a method encompassing a time-consuming multi-PEEP step trial, EIT and esophageal pressure measurements will gain wide clinical acceptance, especially not very early in the course of ventilator treatment or in the operating theater.

However, the PEEP level where transpulmonary driving pressure is lowest can be determined by a rapid two-PEEP-step procedure without both EIT and esophageal pressure measurements [2, 3]. This method, the PEEP step method, is based on the fact that the change in end-expiratory lung volume ($\Delta EELV$) following a PEEP change is determined by the size of the PEEP step ($\Delta PEEP$) and the elastic properties of the lung only, $\Delta EELV = \Delta PEEP \times CL$, i.e., the chest wall does not impede PEEP inflation. (*Details of the determinants of $\Delta EELV$ in e-supplement*). This is an effect of the chest wall striving outwards to a higher volume, 70–80% of total lung capacity during expirations, not only at functional residual capacity, but also at increased EELV and PEEP. If $\Delta EELV$ is measured as the cumulative difference in expiratory tidal volume (VT) between PEEP levels [4], lung compliance can be calculated, $CL_{PSM} = \Delta EELV / \Delta PEEP$

This comment refers to the article available online at <https://doi.org/10.1186/s13054-022-04198-4>.

*Correspondence: ola.stenqvist@aniv.gu.se

Sahlgrenska Academy, Gothenburg University, Gothenburg, Sweden



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

and transpulmonary driving pressure, $\Delta PL = VT/CL_{PSM}$. Also, during PEEP inflation:

- Transpulmonary pressure increases as much as PEEP is increased.
- Transpulmonary driving pressure of a tidal volume equal to $\Delta EELV$ is equal to $\Delta PEEP$.
- Transpulmonary pressure at a certain lung volume is the same irrespective of whether this volume has been reached by tidal inflation or PEEP inflation, i.e., end-inspiratory transpulmonary plateau pressure from low PEEP level is equal to end-expiratory transpulmonary pressure of the high PEEP level.

Because of these features of PEEP inflation, a lung P/V curve can be calculated by a two-PEEP-step procedure from baseline PEEP to end-inspiration at the highest PEEP level. The PEEP level where transpulmonary driving pressure is lowest (least injurious) can then be calculated as the PEEP level half a tidal volume below the steepest point of the lung P/V curve.

Slobod and coworkers did not determine $\Delta EELV$ in their study, but it is possible to illustrate the performance of the PEEP step method by calculating $\Delta EELV$ as $\Delta PEEP \times CL$ between PEEP levels. This enabled the calculation of the complete lung P/V curve and optimal PEEP in the two extreme patients, patients 1 and 3 (Fig. 1). (For details, See Additional file 1: e-supplement.)

There is no safety limit determined for ΔPL , but it can be deduced from the fact that the upper safety limit for ΔPAW is 15 cmH₂O and average ratio of $\Delta PL/\Delta PAW$ is 0.70, which results in a safety limit for ΔPL of $0.7 \times 15 \approx 10$ cmH₂O [2, 5]. Patient 1 has an overall lung compliance of 25 ml/cmH₂O, which is even lower than reported in severe ARDS [5]. Transpulmonary driving pressure at optimal PEEP is almost 10 cmH₂O in spite of a low tidal volume (320 ml). A PEEP increase to 12 cmH₂O would result in a ΔPL of 12.8 cmH₂O and an end-inspiratory transpulmonary plateau pressure of 25 cmH₂O, both dangerously high. In patient 3, with a moderately lowered overall lung compliance, a PEEP increase to 12 cmH₂O only causes a small increase in ΔPL to 6.3 cmH₂O and a transpulmonary plateau pressure of 18 cmH₂O, both well within the safety limits (Fig. 1).

In summary, a complete lung P/V curve and optimal PEEP with lowest transpulmonary driving pressure can be determined by a rapid two-PEEP-step procedure, where $\Delta EELV$ is determined by the ventilator pneumotachograph. Neither esophageal pressure nor EIT is required. The lung P/V curve can be used as clinical decision support to estimate the effect of changes in PEEP and tidal volume on transpulmonary driving and plateau pressure.




Fig. 1 Left panel: Lung P/V curves of patients 1 and 3 with overall lung compliance of 25 and 61 ml/cmH₂O respectively. Overall lung compliance was calculated as end-inspiratory lung volume at the highest PEEP level (12 cmH₂O) divided by end-inspiratory transpulmonary pressure at the highest PEEP level minus 6 cmH₂O (lowest PEEP level). Circles with red filling: end-expiratory transpulmonary pressure (= PEEP). Circles without filling: end-inspiratory transpulmonary pressure. Magenta arrows: tidal lung P/V curves at optimal PEEP obtained by graphical plotting from the lung P/V curve. Data of optimal PEEP tidal volume in italics: VT = tidal volume, ΔPL = transpulmonary driving pressure, CL = lung compliance, PEEP_{opt} = optimal PEEP level, PL_{plat} = end-inspiratory transpulmonary pressure. Right panel: Light gray lung P/V curves of patients 1 and 3 can be used as clinical decision support as a tidal lung P/V curve with any combination of PEEP and VT will be positioned on the complete lung P/V curve. This makes it possible to estimate the effect of changes in PEEP and tidal volume. In this case, the tidal lung P/V curves (black arrows) are depicted starting from PEEP 12 cmH₂O. In patient 3 with a moderately decreased overall lung compliance to 61 ml/cmH₂O ("normal" lung compliance 90–110 ml/cmH₂O [3]), transpulmonary driving pressure increases marginally and remains within safe limit, i.e., the patient responds favorably to PEEP. In patient 1, an increase in PEEP to 12 cmH₂O results in a ΔPL 40% higher almost 13 cmH₂O than at PEEP 9 cmH₂O with significant risk of ventilation induced lung injury

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13054-022-04240-5>.

Additional file 1. E-supplement.

Author contributions

OS wrote the manuscript. The author read and approved the final manuscript.

Funding

The author declares that there is no funding.

Availability of data and materials

Not applicable.

Declarations

Ethics approval and consent to participate

Not applicable.

Competing interests

OS holds shares in the Lung Barometry Sweden Company.

Received: 8 November 2022 Accepted: 12 November 2022

Published online: 29 November 2022

References

1. Slobod D, Leali M, Spinelli E, Grieco DL, Spadaro S, Mauri T. Integrating electrical impedance tomography and transpulmonary pressure monitoring to personalize PEEP in hypoxemic patients undergoing pressure support ventilation. *Crit Care*. 2022;26(1):314.
2. Grivans C, Stenqvist O. Gas distribution by EIT during PEEP inflation: PEEP response and optimal PEEP with lowest trans-pulmonary driving pressure can be determined without esophageal pressure during a rapid PEEP trial in patients with acute respiratory failure. *Physiol Meas*; 2022.
3. Persson P, Stenqvist O. Protective positive end-expiratory pressure and tidal volume adapted to lung compliance determined by a rapid positive end-expiratory pressure-step procedure in the operating theatre: a post hoc analysis. *Br J Anaesth*. 2022;128(4):e284–6.
4. Grivans C, Lundin S, Stenqvist O, Lindgren S. Positive end-expiratory pressure-induced changes in end-expiratory lung volume measured by spirometry and electric impedance tomography. *Acta Anaesthesiol Scand*. 2011;55(9):1068–77.
5. Gattinoni L, Pelosi P, Suter PM, Pedoto A, Vercesi P, Lissoni A. Acute respiratory distress syndrome caused by pulmonary and extrapulmonary disease: different syndromes? *Am J Respir Crit Care Med*. 1998;158(1):3–11.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

