

MATTERS ARISING

Open Access



Serial assessments of cardiac output and mixed venous oxygen saturation in comatose patients after out-of-hospital cardiac arrest

Tobias Zimmermann^{1*}, Pedro Lopez-Ayala² and Mervyn Singer¹

We read with great interest the article “Serial assessments of cardiac output and mixed venous oxygen saturation in comatose patients after out-of-hospital cardiac arrest” by Grand et al. [1] While the reported findings are generally in line with what we would predict from a physiological point of view, we query the rigor of the statistical analyses.

To illustrate the relationship between mortality and hemodynamic variables, the authors used a Cox proportional hazards model with smoothing splines to attain the predicted values. It is unclear whether this was an univariable or multivariable model. The authors then present Figure 4 of their paper as a plot of the “Hazard ratio of mortality as a function of first measured mixed venous oxygen saturation [...] and first measured cardiac index [...] values during intensive care after cardiac arrest.” However, since no reference point was chosen, this represents a (relative) hazard plot and not a hazard ratio plot [2]. A dose–response plot with an appropriate reference point (illustrated in Fig. 1), defined based on prior clinical

knowledge, should be presented instead. The literature offers examples of such hazard ratio dose–response plots [3]. This could potentially affect the conclusion that can be drawn; for example, depending on the location of the chosen reference point, a significant association between cardiac index and mortality might be revealed.

Could the authors also elaborate on their choice of relaxing linearity assumptions with splines for graphical illustration but not for the regression models? Seeing the nonlinear relationship between the variables of interest and the outcome graphically represented in Figure 4 of their paper, should this not be taken into account when fitting the multivariable prediction model? Categorizing cardiac index into quarters is a suboptimal way of dealing with nonlinearity. This should be avoided [4] especially as splines have already been introduced into the analysis that are capable of relaxing linearity assumptions within prediction models [2, 5]. This could have important implications on the central conclusions drawn from the analysis.

This comment refers to the article available online at <https://doi.org/10.1186/s13054-023-04704-2>.

*Correspondence:

Tobias Zimmermann
tobias.zimmermann@nhs.net

¹ Bloomsbury Institute of Intensive Care Medicine, University College London, London, UK

² Cardiovascular Research Institute Basel, University Hospital Basel, University of Basel, Basel, Switzerland



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.




Fig. 1 Example of a dose–response hazard ratio plot based on the open access Mayo Clinic Primary Biliary Cholangitis Dataset. A multivariable Cox proportional hazards model was fitted, conditioning on treatment received and bilirubin level. To relax the linearity assumption, restricted cubic splines with four knots were fitted for age and bilirubin. Note the selected reference point where the 95% CI collapses. The R code for creating this figure can be found online (https://gitlab.com/tobiaszimmermann/lte_critcare_grand_2023)

Author contributions

TZ and PL-A wrote the manuscript under the guidance of MS. All authors reviewed the manuscript.

Funding

Not applicable.

Availability of data and materials

The Mayo Clinic Primary Biliary Cholangitis Dataset used to create the example of a dose–response hazard ratio plot is part of the R “survival” package (<https://rdrr.io/cran/survival/man/pbc.html>) [6]. The R code for creating this figure can be found online (https://gitlab.com/tobiaszimmermann/lte_critcare_grand_2023).

Declarations

Ethics approval and consent to participate

Not applicable.

Competing interests

The authors declare no competing interests.

Received: 3 November 2023 Accepted: 10 November 2023

Published online: 20 November 2023

References

- Grand J, Hassager C, Schmidt H, et al. Serial assessments of cardiac output and mixed venous oxygen saturation in comatose patients after out-of-hospital cardiac arrest. *Crit Care*. 2023;27:410.

- Harrell FE. *Regression modelling strategies*. 2nd ed. New York: Springer; 2015.
- Crane PK, Walker R, Hubbard RA, et al. Glucose levels and risk of dementia. *N Engl J Med*. 2013;369:540–8.
- Naggara O, Raymond J, Guilbert F, et al. Analysis by categorizing or dichotomizing continuous variables is inadvisable: an example from the natural history of unruptured aneurysms. *AJNR Am J Neuroradiol*. 2011;32:437–40.
- Ma J, Dhiman P, Qi C, et al. Poor handling of continuous predictors in clinical prediction models using logistic regression: a systematic review. *J Clin Epidemiol*. 2023;161:140–51.
- Therneau TM, Grambsch PM. *Modeling survival data: extending the cox model*. New York: Springer; 2013.

Publisher’s Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.