

RESEARCH

Open Access



# Exploring the effectiveness of eHealth interventions in treating Post Intensive Care Syndrome (PICS) outcomes: a systematic review

Daniel Jie Lai<sup>1,2</sup>, Zhao Liu<sup>3,4</sup>, Elaine Johnston<sup>5,6</sup>, Lisa Dikomitis<sup>6,7</sup>, Teresa D'Oliveira<sup>1,8†</sup> and Sukhi Shergill<sup>6,8\*†</sup>

## Abstract

**Background** It remains unclear how to optimise critical care rehabilitation to reduce the constellation of long-term physical, psychological and cognitive impairments known as Post Intensive Care Syndrome (PICS). Possible reasons for poor recovery include access to care and delayed treatment. eHealth could potentially aid in increasing access and providing consistent care remotely. Our review aimed to evaluate the effectiveness of eHealth interventions on PICS outcomes.

**Methods** Studies reporting eHealth interventions targeting Post Intensive Care Syndrome outcomes, published in Medline, CINAHL, PsycINFO, Embase, and Scopus from 30th January 2010 to 12th February 2024, were included in the review. Study eligibility was assessed by two reviewers with any disagreements discussed between them or resolved by a third reviewer. Study quality and risk of bias were assessed using the Mixed Method Appraisal Tool. Further to the identification of effective strategies, our review also aimed to clarify the timeline of recovery considered and the outcomes or domains targeted by the interventions.

**Results** Thirteen studies were included in our review. Study duration, eHealth intervention delivery format, and outcome measures varied considerably. No studies reported a theory of behavioural change and only one study was co-produced with patients or carers. Most studies were conducted in the early post-discharge phase (i.e., < 3 months) and had feasibility as a primary outcome. The cognitive domain was the least targeted and no intervention targeted all three domains. Interventions targeting the psychological domain suggest generally positive effects. However, results were underpowered and preliminary. Though all studies were concluded to be feasible, most studies did not assess acceptability. In studies that did assess acceptability, the main facilitators of acceptability were usability and perceived usefulness, and the main barrier was sensitivity to mental health and cognitive issues.

**Conclusion** Our systematic review highlighted the promising contributions of eHealth with preliminary support for the feasibility of interventions in the early stages of post-critical care rehabilitation. Future research should focus on demonstrating effectiveness, acceptability, the cognitive domain, and multi-component interventions.

**Keywords** Critical care, Critical Illness, Critical care rehabilitation, Post-intensive care syndrome, EHealth, Digital health technologies

<sup>†</sup>Teresa D'Oliveira and Sukhi Shergill these authors are joint senior authors on this work.

\*Correspondence:

Sukhi Shergill

sukhi.shergill@kcl.ac.uk

Full list of author information is available at the end of the article



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## Background

Post Intensive Care Syndrome (PICS) has been increasingly recognised as an urgent problem among critical care survivors [1–4]. This is characterised as a sequelae of new or worsened physical, psychological, and cognitive impairments after critical illness which has significant impacts on functional outcomes, Health-Related Quality of Life (HRQoL), and employment [5–8]. The establishment of a rehabilitation pathway is essential for successful PICS management.

Critical care rehabilitation consists of four phases: acute recovery and prevention within the critical care unit, recovery in the hospital ward, the first 3 months after hospital discharge termed the early post-discharge period, and the late post-discharge period which can span years after discharge [9]. Our review terms the three phases after critical care discharge as the ‘post-critical care’ phases. The effectiveness of current interventions in the post-critical care phases are limited with most targeting the late post-discharge period [10, 11]. This limited effectiveness could be due to the time points chosen to begin rehabilitation (i.e., a later start of rehabilitation). The early post-discharge period is deemed a crucial recovery point where critical care survivors are most vulnerable. These impacts are further magnified by regional health inequalities that restrict access to care [12]. There is a need for earlier intervention and continuity of care.

The use of electronic Health (eHealth) is presented by the literature as a solution to minimise health inequalities and facilitate earlier intervention. eHealth technologies are characterised by 1) enabling the storage, retrieval, and transmission of data, 2) supporting clinical decision-making, and 3) facilitating remote care [13]. These technologies include mobile applications, video conferencing, virtual reality, web platforms and wearable technology. The use of eHealth has proliferated within critical care. For example, the tele-critical care model aids in addressing workforce shortages, provides better access to specialist expertise, reduces patient transfers, and lowers ICU mortality [14, 15]. However, efforts in harnessing the benefits of eHealth have only just begun in post-critical care.

Studies conducted in the last 3 years demonstrate a demand for tools that can detect and measure rehabilitation of PICS symptoms [16]. The use of eHealth interventions to rehabilitate patients in the early post-discharge phase could promote better PICS recovery. Evidence from other chronic patient populations like heart failure, stroke and diabetes has shown promising results in eHealth’s effectiveness on post-hospital disease management, medicine adherence, and health-related quality of life [17–19]. However, specific identification and evaluation in a post-critical care setting has yet to be done. To

our knowledge, this is the first comprehensive review of eHealth’s impact on PICS outcomes during the critical care rehabilitation phase. This encompasses the in-hospital, early, and late post-discharge phases. The objective of the review is to identify effective strategies using eHealth that target PICS, their timeline in the recovery path and the outcomes addressed. As primary outcomes, we consider the PICS domains (physical, psychological and cognitive) targeted by the eHealth interventions, the recovery phase these interventions are implemented and their effectiveness. Secondary objectives include the feasibility of these eHealth interventions, acceptability, and identification of the barriers, and facilitators of eHealth intervention uptake.

## Methods

This systematic review is reported based on the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) [20]. The study was registered and published in the International Prospective Register of Systematic Review databases (PROSPERO registration number: CRD42023463036) [21]

### Search strategy and selection criteria

#### Search strategy

The following databases were searched: Medline, CINAHL, PsycINFO, Embase, and Scopus. Reference lists from key articles were also checked for any additional articles that fit the inclusion criteria. Due to the rapid innovation of eHealth technologies, studies that were published from 30th January 2010 to 12th February 2024 were included in this review. No restrictions were imposed on the language of publication.

The PICO framework [22] was used to identify key terms and develop the search string. PIO was used as there was no restriction imposed on the study design. The comparator category was not included in the search strategy to expand the articles picked up. The categories were defined as (P): Post Intensive Care patients; (I): eHealth interventions; (O) Post Intensive Care Syndrome outcomes (Physical, Psychological, Cognitive). The search string was tailored to fit the querying format of each database and can be found in Supplementary Material S1.

#### Study inclusion and exclusion criteria

Eligible studies included i) adults over the age of 18 who have been discharged from critical care (in the hospital ward, early post-discharge, and late post-discharge), ii) the inclusion of one or more eHealth interventions implemented in any of the three phases of post-critical care recovery, iii) PICS domains were measured as an outcome, vi) full text published in peer reviewed journals. There were no restrictions made on the study design

and the language of publications. As current eHealth definitions proposed in the literature are very broad and general, we operationalised what constitutes an eHealth intervention using the definition by Black et al., [13] which was conceptualised to aid the categorisation of eHealth interventions using themes generated from 53 systematic reviews. The eHealth inclusion and exclusion criteria were developed based on this definition and the types of eHealth interventions were categorised in these categories.

- (1) Telemedicine
- (2) Telerehabilitation
- (3) Self-directed interventions
- (4) Remote patient monitoring (wearables, sensors)
- (5) Virtual Reality (VR)

Studies excluded consisted of (i) no evidence of eHealth intervention, (ii) Paediatric (children) ICU, (iii) neonatal/prenatal ICU, (iv) systematic reviews and meta-analyses, (v) conference abstracts, and (vi) study protocols.

#### **Selection process**

Two reviewers (DL, ZL) independently screened the articles according to the stipulated inclusion and exclusion criteria. During the titles and abstract screening stage, screening procedures proposed by Adams et al. [23] were used. The first reviewer (DL) screened all titles and abstracts, while the second reviewer (ZL) screened a 10% random selection of articles. There was substantial inter-rater reliability between the reviewers (Kappa = 0.66; percentage agreement = 98.8%). Full-text screening was done independently by DL and ZL with almost perfect agreement (Kappa = 0.95, percentage agreement = 98.3%). Any disagreements were discussed between the two reviewers until a consensus was reached. When consensus could not be reached, the dispute was solved with the consultation of a senior team member (TD).

#### **Data extraction**

Data extracted consisted of study characteristics (Author/year; Country; Study design; Population; Post-critical care timepoint; Sample size/Control (if any); Study duration), eHealth intervention characteristics (Intervention; Type of eHealth intervention; Delivery Format; Outcome Measures; Findings).

Feasibility was measured and assessed in different ways due to the variation of eHealth interventions. Feasibility data extracted included the feasibility outcome defined by authors, attrition, definition of intervention adherence, adherence rate, reasons for participant withdrawal, and author's conclusions.

Data extracted for acceptability consisted of how acceptability was assessed (acceptability measure), main findings, and reported barriers and facilitators in intervention uptake. Data extraction was done in duplicate by two reviewers (DL and ZL) who worked independently.

#### **Risk of bias and quality assessment**

Two reviewers (DL, and ZL) independently assessed the risk of bias and the quality of studies using the Mixed Methods Appraisal Tool (MMAT) [24]. The tool has 5 quality criteria examining and evaluating the appropriateness of a study's aims, methodology, design, data collection, data analysis, presentation of findings, discussion, and conclusion. The quality criteria are rated with 'Yes', 'No', or 'Can't tell' and are evaluated based on study design. Criteria for a randomised controlled trial are different from a non-randomised trial (quality criteria can be found in Supplementary Material S2). Each study was scored using percentages based on the recommendations by Pace et al. [25]. Any disagreements were resolved through discussion between the two reviewers.

#### **Data synthesis and analysis**


A quantitative analysis of outcomes or meta-analysis could not be done due to the heterogeneity of the study designs, outcome measures used, eHealth interventions, and the critical care population. With the included studies, a qualitative narrative synthesis was undertaken to summarise the primary and secondary outcomes of interest. Data were grouped based on the main outcomes listed in the data extraction section.

#### **Results**

Initial database searches yielded 3,673 articles. The deduplication of 428 articles led to a total of 3,245 titles and abstracts screened. In accordance with the exclusion criteria, 3,186 articles were excluded leaving 59 articles for full-text retrieval. Out of the 59 articles, 13 met the inclusion criteria for the current review. Figure 1 presents the PRISMA diagram documenting the processes of identifying, screening, and selecting included papers.

#### **Study characteristics**

Studies were conducted across 7 countries with the majority coming from the United States (6/13). A total of 548 participants were enrolled across 13 studies. The sample sizes ranged from 5 to 89 with participant ages ranging from 47 to 72 years. Study design varied considerably across the studies with 46% (6/13) of studies being Randomised Controlled feasibility Trials (RCT) [31–33, 36, 41, 42], 38.4% (5/13) prospective observational cohort studies [34, 35, 38, 40, 43], and 15.3% (2/13) qualitative studies [37, 39]. None of the studies reported



**Fig. 1** PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) flow diagram documenting the processes of identification, screening, and article inclusion. Latest search 12th February 2024

any underpinning theory of behaviour change and only 1 study [40] reported co-production efforts during intervention development. The characteristics and intervention descriptions of included studies are summarised in Table 1.

**Interventions targeting PICS**

There was a wide range of different eHealth interventions and delivery formats. 3 studies investigated telerehabilitation [27, 31, 33], 2 studies investigated telemedicine [26, 35], 2 studies investigated patient monitoring [29, 30], 3 studies investigated virtual reality

[36–38], and 1 study investigated a self-directed eHealth intervention [28].

Out of the three domains, eHealth interventions targeted the psychological domain most frequently [26, 28, 33, 35–37], followed by the physical domain [26, 27, 29–31] and the cognitive domain being the least targeted [27, 31, 38]. Only three study teams designed interventions that covered two PICS domains [26, 27, 31]. There were no eHealth interventions that targeted all three PICS domains in tandem. Table 2 summarises the relationship between the intervention delivery format and the domains targeted.

**Table 1** Study characteristics, description of eHealth interventions and main findings

Author, Year	Country	Study Design	Population	Time point	Sample/Control	Duration	Intervention	Type of ehealth intervention	Delivery format	Outcome Measures	Main Findings
Balakrishnan et al. [26]	US	Feasibility RCT	Covid-19 patients	Early post-discharge	40/TAU <sup>§</sup>	10 weeks	Telemedicine visits 14 days post-hospital discharge in Covid-19 and ARDS survivors	Telemedicine/ Video call	Video call	6MWT EQ5D5L	No significant difference in HFQOL and anxiety
Capin et al. [27]	US	Feasibility RCT	Covid-19 patients	Early post-discharge	44/Active*	12 weeks	Telerehabilitation program which included 12 exercise sessions led by a physiotherapist	Tele-rehabilitation	App	30 s Chair stand test TUG Step count MoCA	No significant difference in physical outcomes No significant difference in cognitive outcomes
Cox et al. [28]	US	Feasibility RCT	Medical, cardiac, and surgical ICU	Early post-discharge	80/Active*	12 weeks	Self-directed mindfulness application which delivered 4 mindfulness sessions consisting of background videos, guided meditation and educational materials	Self-Directed eHealth intervention	App	PHQ-15 PTSS PHQ-9 GAD-7 EQ5D5L	Mindfulness group has lower PTSD and Depressive symptoms and anxiety scores compared to group that received education Feasibility, acceptability, and usability concluded

**Table 1** (continued)

Author, Year	Country	Study Design	Population	Time point	Sample/Control	Duration	Intervention	Type of eHealth intervention	Delivery format	Outcome Measures	Main Findings
Denehy et al. [29]	Australia	Observational Quasi-experimental	Medical ICU	In-hospital	53	1 week	Critical care survivors wore an actigraph for 7 days at 2 months post-hospital discharge	Remote patient monitoring	Wearable	6MWT TUG SF-36 Step count	Absence of chronic disease was significantly associated with increased distance walked ( $p < .000$ ) where chronic disease explained 33.5% of the variance in average distance walked ICU survivors were inactive when quantitatively measured at 2 months after hospital discharge 63% of participants did not reach 30 min of moderate physical activity

**Table 1** (continued)

Author, Year	Country	Study Design	Population	Time point	Sample/Control	Duration	Intervention	Type of eHealth intervention	Delivery format	Outcome Measures	Main Findings
Estrup et al. [30]	Denmark	Observational Quasi-experimental	Medical, surgical ICU	In-hospital	44	1 week	Activity levels measured for 7 days in-hospital. Physical function outcomes were measured 3 months post-hospital discharge	Continuous patient monitoring	Wearable	CPAx Step count	Improved physical function in most patients 3-month post-hospital discharge Physical function correlated with mean daily activity ( $p=0.017$ ), maximum activity on second day ( $p=.0053$ ), and total activity in the day-time ( $p=0.0058$ )
Jackson et al. [31]	US	Feasibility RCT	Medical, surgical ICU	Early post-discharge	21/TAU <sup>s</sup>	12 weeks	A hybrid cognitive, physical and functional rehabilitation, physical and functional rehabilitation sessions were conducted via video call. Cognitive rehabilitation was done in-person	Tele-rehabilitation	Video call	TUG Tower test MMSE	Improvements in cognitive outcomes Improvements in instrumental activities of daily living (driving, shopping)
Kovaleva et al. [32]	US	Qualitative	ICU Sepsis and/or ARDS diagnosis	Late post-discharge	24	12 weeks	Video conference at 3- and 12-weeks post-hospital discharge	Telemedicine	Video call	Patient experience	All participants found telemedicine visit acceptable Technology was easy to use

**Table 1** (continued)

Author, Year	Country	Study Design	Population	Time point	Sample/Control	Duration	Intervention	Type of eHealth intervention	Delivery format	Outcome Measures	Main Findings
Park et al. [33]	US	Observational Quasi-experimental	Covid-19	Late post-discharge	18	Up to 14 sessions	Up to 14 virtual, evidence-based psychotherapy sessions for patients. Intervention integrated psychoeducation, cognitive restructuring, acceptance, exposure-based, and mindfulness	Tele-psychotherapy	Video call	PHQ-9 GAD-7	Significant decrease in participants meeting anxiety (57.1% decrease) and depression (57.2% decrease)
Parker et al. [34]	Australia	Qualitative	ARDS survivors	Late post-discharge	10	N. A	Application which consisted of educational resources on rehabilitation, a mood tracker, and a goal tracker	Educational	App	Acceptability	80% stated application was easy to use 37% stated it would be helpful to have an application demonstration Median score for app 'helpfulness' was the maximum score of 5, IQR [4.24–5.00]
Rose et al. [35]	UK	Observational Quasi-experimental	Medical, Surgical ICU	Early post-discharge	5	12 weeks	Delivered through online/mobile platform. The platform provided assessments of baseline status and recovery barriers, tailored e-resources based on recovery barriers, patient recovery e-diary	Telemedicine	Web-based/ App	Short-term goal completion	Two patients reported achieving short-term goals, 2 partially achieved goals, and 1 did not achieve



**Table 1** (continued)

Author, Year	Country	Study Design	Population	Time point	Sample/Control	Duration	Intervention	Type of eHealth intervention	Delivery format	Outcome Measures	Main Findings
Vlake et al. [36]	Netherlands	Multi-centre feasibility RCT	ICU Sepsis or septic shock diagnosis	In-hospital	50/Active*	1 session	Patients in the step-down ward go through the ICU-VR intervention which includes different scenes in the ICU unit. Patients were then followed up 2 days, 1 week, 1 month, and 6 months after VR intervention	Virtual Reality intervention	Virtual Reality headset	IES-R BDI-II MCS-12 EQ5D5L	PTSD and depression reduced in VR intervention group 2 days after exposure and persisted throughout follow up time-points HRQoL improved in VR intervention group up to 1 month after VR intervention. Did not improve 6 months after intervention VR intervention group experienced greater sense of presence, involvement and experienced realism
Vlake et al. [37]	Netherlands	Multi-centre feasibility RCT	Covid-19	Late post-discharge	89/TAU <sup>§</sup>	1 session	VR intervention is the same as intervention above but was trialed on a post-hospital discharge sample. Participants were followed up 1 and 3 months after 3-month post-ICU clinic appointment	Virtual Reality intervention	Virtual Reality headset	IES-R HADS SF-36 EQ5D5L	Psychological outcomes and quality of life did not improve Satisfaction in ICU aftercare rated higher in intervention group

**Table 1** (continued)

Author, Year	Country	Study Design	Population	Time point	Sample/Control	Duration	Intervention	Type of eHealth intervention	Delivery format	Outcome Measures	Main Findings
Wood et al. [38]	Canada	Observational Quasi-experimental	Medical, Surgical ICU	Late post-discharge	70	2 sessions	A screening tool where participants completed 7 tasks within a VR environment	Virtual Reality Screening Tool	Virtual Reality	RBANS KINARM (VR Intervention)	Prevalence of cognitive and sensorimotor impairments 9/28 (32%) and 3/22 (14%) participants displayed global cognitive impairment Screening tool caused little to no procedural discomfort

None of the included studies reported an explicit behavioural theory or model of change

RCT Randomised Controlled Trial, TAU/Treatment as usual, HRQoL Health Related Quality of Life, PTSD Post Traumatic Stress Disorder, ICU Intensive Care Unit, M Mean, SD Standard Deviation, ARDS Acute Respiratory Distress Syndrome, VR Virtual Reality

Outcome Measure Abbreviations: 6MWT 6 Minute Walk Test, EQ5D5L 5 Europe Quality of Life 5 Dimensions 5 Levels, TUG Timed Up and Go Test, MoCA Montreal Cognitive Assessment, PHQ-9/15 Patient Health Questionnaire (9 or 15 question version), GAD-7 Generalised Anxiety Disorder – 7 questions, PTSD Post Traumatic Stress Disorder – 7 questions, PDS Post Traumatic Syndrome Scale, SF-36 Short Form-36 questions, CPAX Chelsea Critical Care Physical Assessment Tool, MMSE Mini-Mental State Examination, IES-R Impact of Event Scale- Revised, BDI-II Beck Depression Inventory 2, MCS-12 Mental health Component Summary-12 questions, HADS Hospital Anxiety and Depression Scale, RBANS Repeatable Battery for the Assessment of Neuropsychological Scale

<sup>†</sup> Rose et al., (2021) was the only included study which reported to have done co-production and patient and public involvement work during the eHealth intervention phase

<sup>§</sup> Treatment as Usual control groups underwent usual outpatient clinics

\* Active control groups were provided education materials and a face-to-face equivalent of the eHealth intervention

**Table 2** Summary of targeted PICS domains of each eHealth intervention

Author/Year	Intervention delivery format	Physical	Psychological	Cognitive
Denehy et al. [29]	Wearable sensor	x		
Estrup et al. [30]	Wearable sensor	x		
Jackson et al. [31]	Video conference	x		x
Capin et al. [27]	Video conference	x		x
Balakrishnan et al. [26]	Video conference	x	x	
Park et al. [33]	Video conference		x	
Cox et al. [28]	Application		x	
Rose et al. [35]	Web/application		x	
Vlake et al. [36]	Virtual reality		x	
Vlake et al. [37]	Virtual reality		x	
Wood et al. [38]	Virtual reality			x

### Timing of interventions

Most of the included studies (5/11 studies) chose the early post discharge phase [26–28, 31, 35]. Three studies [29, 30, 36] were conducted in-hospital and 3 studies during the late post-discharge [33, 37, 38].

### eHealth intervention effects on PICS outcomes

#### Outcome measures

There were a variety of outcome measures for each PICS domain. Physical measures include 6MWT [26, 29], TUG [27, 29, 31], CPAx [30], actigraphy step count [29, 30], and 30-s chair stand [27]. Psychological outcome measures included the HADS [37], PHQ [28, 33], GAD-7 [28, 33], BDI-II [36], SF-36 [29], MCS-12 [36], PTSS [28] and IES-R [36, 37]. Cognitive measures included MoCA [27], MMSE [31], RBANS [38]. Studies measuring Health-Related Quality of Life all used the EQ-5D-5L [28, 36, 37].

#### Physical outcomes

The impact of eHealth interventions on physical function was mixed. Whilst Jackson et al. [31] found a significant effect on physical function with a multi-component telerehabilitation, Capin et al. [27] did not find any significant effects on physical function with a tele-physical therapy intervention. A significant improvement in physical function at 3 months post-discharge was significantly correlated with mean daily activity [30]. An absence of chronic disease is a majorly significant ( $p < 0.000$ ) predictor of increased distance walked post-hospital discharge explaining 33.5% of the variance in mean distance walked [29].

#### Psychological and cognitive outcomes

Of the 6 studies that targeted psychological outcomes, 4 studies showed significant reductions in anxiety [33], depression [28, 33, 36], and Post Traumatic Stress

Disorder [28, 36]. Only 2 studies showed no effects [26, 37].

Two studies that targeted cognitive outcomes used the same telerehabilitation programmes used in the physical outcomes section [27, 31]. Capin et al. [27] did not find any improvement in cognitive outcomes while Jackson et al. [31] found significant improvement in executive functioning. Wood et al. [38] tested a cognitive screening tool and found less pronounced cognitive impairment 12 months after hospital discharge.

### Secondary outcomes

#### Feasibility

All the included studies which explored feasibility (9 out of 13 studies) demonstrated the feasibility of the various eHealth interventions. Outcome measures used to evaluate feasibility varied. All studies used adherence as an outcome of feasibility. Other outcomes include Attrition [28], safety through reported adverse events [27], VR immersion, and motion sickness [36]. All studies had an adherence rate of more than 70%. One study had 71% adherence [28], 4 studies had >75% adherence [29–31, 33], 1 study had 83% adherence [27], 1 study had 90% adherence [26], and 2 studies had 100% adherence [36, 37]. A summary of the defined feasibility outcomes and findings is summarised in Additional file 1: Table S3.

#### Acceptability of eHealth interventions

Studies which reported acceptability (5 out of 13) included two qualitative studies [32, 34] and 3 RCTs [26, 27, 37]. Acceptability measures mainly evaluated participant satisfaction and perceptions of the intervention. All studies concluded the intervention to be acceptable. The 3 RCT studies evaluated acceptability using a questionnaire and reported high participant satisfaction.

The two qualitative studies focused on the experiences of a telemedicine intervention and an app-based mood monitoring prototype system [32, 34]. Both studies assessed acceptability through semi-structured interviews and reported barriers and facilitators in intervention uptake.

Most themes considered the sensitivity of mental health and cognitive issues as barriers. Participants from Kovaleva et al. [32] study mentioned that neuropsychological assessments felt ‘embarrassing’ when other clinicians were present in the video call while participants in Parker et al. [34] study thought ‘depression’ was too stigmatising and suggested the term emotions/states as an option.

Usability and perceived usefulness were identified as the main facilitators of the use of eHealth interventions. Facilitators in the acceptability of eHealth interventions included the ease of using the intervention platforms, the convenience, and viewing the platform as a motivator of recovery. A summary of all the acceptability findings can be found in Additional file 1: Table S4.

#### Quality assessment and risk of bias of included studies

Quality assessments used the MMAT tool [24] with most studies running quantitative randomised controlled trials. Though included RCTs varied in quality, most of the RCT studies were of high quality with 4 of 6 studies scoring 80% [27, 28, 31, 37] and 2 studies were of moderate quality scoring 60% [26, 36]. The main limitations impacting study quality were due to incomplete outcome data and the inability to ‘blind’ participants. There was

a greater variance in study quality for non-randomised quantitative studies with 2 high-quality studies scoring 80% [29, 30], 2 studies moderate quality studies scoring 60% [33, 38] and 1 low quality study scoring 20% [35]. The main limitations that impacted the low-quality study were the representativeness of the sample, selection of measures, and incomplete description of intervention as intended. The two qualitative studies were high-quality at 80% [32] and 100% [34]. The detailed rating and scoring of the MMAT tool can be found in Table 3

#### Discussion

The main objectives of the study were to systematically assess and explore eHealth’s effectiveness in alleviating PICS impairments, when in the recovery path these are implemented, and the domains being targeted by each intervention. There was a great variety of eHealth interventions with most studies focussing on the physical and psychological domains. Most studies were conducted in the early post-discharge phase and had feasibility as a primary outcome. There is great heterogeneity in the outcome measures used to assess PICS domains, feasibility and acceptability. Nevertheless, findings from the review suggest that eHealth interventions are feasible in a post-critical care setting with further research required in measuring effectiveness.

Though there is variation in the outcome measures used to assess PICS outcomes, the majority of the studies used measures recommended by published core outcome sets (COS). The lack of consistency is due to the different COS available. Remote physiotherapy interventions used

**Table 3** Mixed method appraisal tool risk or bias rating scores

Author (Year), Country	1.1	1.2	1.3	1.4	1.5	2.1	2.2	2.3	2.4	2.5	3.1	3.2	3.3	3.4	3.5	Score(%)
Qualitative																
Kovaleva et al. [32]	Y	N	Y	Y	Y											80%
Parker et al. [34]	Y	Y	Y	Y	Y											100%
Quantitative Randomised																
Balakrishnan et al. [26]						Y	Y	N	N	Y						60%
Capin et al. [27]						Y	Y	Y	N	Y						80%
Cox et al. [28]						Y	Y	Y	Y	N						80%
Jackson et al. [31]						Y	Y	Y	Y	N						80%
Vlake et al. [36]						Y	Y	Y	N	N						60%
Vlake et al. [37]						Y	Y	Y	N	Y						80%
Quantitative Non-randomised																
Denehy et al. [29]											Y	Y	Y	N	Y	80%
Estrup et al. [30]											Y	Y	Y	N	Y	80%
Park et al. [33]											Y	N	Y	N	Y	60%
Rose et al. [35]											N	Y	N	U	U	20%
Wood et al. [38]											Y	Y	N	N	Y	60%

Y Yes, N No, U Unknown/Can't Tell

a COS focussing on critical care physical rehabilitation [39], while other interventions used a mixture of clinically based COS [40] and COS for clinical research [41]. COS is produced to reduce outcome measure heterogeneity and enable better data synthesis [42]. However, none of the studies reported which outcome sets the measures were selected from. To meet the aims of producing a COS, future studies should report how measures were chosen and identify if a specific COS was used. This will provide consistency in reporting and ease for researchers to compare results across eHealth interventions.

The effects of the eHealth interventions on PICS outcomes were mixed. This is the case for physical and psychological outcomes. The majority of studies targeting psychological outcomes had more interventions reporting positive effects. Vlasek et al. [37] did not find significant improvements in psychological outcomes in a late post-discharge sample. However, a prior study conducted by the same authors found an improvement in psychological outcomes in an in-hospital sample that persisted across other follow-up time points [36]. Prior systematic reviews on post-critical care rehabilitation have highlighted the importance of intervention timing [11, 43]. Just as early mobilisation in the critical care ward can alleviate the risk of PICS development [44], there may be an optimal window across the post-critical care recovery path for certain interventions to be effective.

Cognitive outcomes were the least targeted out of the three PICS domains. Studies investigating this outcome observed improvement with multi-component rehabilitation, Jackson et al. [31] attributed significant effects in physical and cognitive outcomes when combining rehabilitation of the two domains together, a result that contrasts with Capin et al. [27] programme which focussed on physical function only. The potential benefits and synergistic effects of performing physical exercise and cognitive training have been documented in other populations [45]. Interrelationships among the three domains are presented through the prevalence of PICS symptom comorbidities. Heesaker et al. [46] observed that mental health and cognitive impairment always occur simultaneously with the other two domains. Marra et al. [47] reported a combination of mental health and cognitive impairment occurring more frequently than other combinations. Kang et al. [48] built on those studies and found that 41.1% of critical care survivors with PICS had symptoms in two or more domains with Physical-Mental symptoms being the most prevalent. With these potential effects, the review found that there has yet to be an intervention that targets these three domains. The incorporation of the cognitive domain is still incipient, and more evidence is required to determine the impact of multi-component interventions.

None of the included studies reported on a theory of behaviour change and only one study [35] reported evidence of co-producing the intervention. Recent guidelines from the Medical Research Council recommend complex health interventions to be co-produced and underpinned by the behavioural theory of change as it increases the effectiveness of behaviour change [49–51]. There is a possibility that behavioural theories have been implied and not discussed explicitly. Goal setting was used in the digital pathway intervention by Rose et al., [35], app-based Mindfulness [28] and tele-psychotherapy [33] rely on the mechanisms of change brought by the therapeutic approaches. Nevertheless, explicit reporting of theories used as well as evidence of co-production is integral in evaluating complex health interventions.

Most studies point to the feasibility of implementing eHealth interventions. With regards to acceptability, studies that assessed it deemed the eHealth interventions feasible. The implementation of eHealth interventions into day-to-day clinical practice has been challenging [52]. The decision to adopt an eHealth intervention requires careful management of both patient and staff expectations [53]. Clinicians and hospital staff need to believe that the intervention can improve care and efficiency. They need to be on board, involved, and receive consistent support during the adoption [54]. The success of eHealth implementation is also determined by patient engagement and uptake. This is especially challenging in older patient populations like critical care survivors. The themes of usability and perceived usefulness highlighted in this review were in line with older patients with chronic conditions [55], older patients with cancer [56], and the general older population [56, 57]. Critical care survivors were more likely to adhere to eHealth interventions when they are easy to use, convenient and perceived as a motivator towards recovery. The continuous contact between patients and the clinical team through telemedicine visits supported the perceptions of care continuance, thus increasing the perceived usefulness and adherence to eHealth interventions. Despite the alignment with research on senior populations, acceptability was only assessed by 5 out of 13 studies which limits the generalisability of findings in a post-critical care population. Further research is needed to address the specific barriers and facilitators for eHealth uptake and engagement in this population.

#### **Study limitations**

One limitation of this review is the infancy of the current research area. The primary objective of studies included in the review was to assess the feasibility of the intervention resulting in underpowered studies with small samples. The effects of eHealth on each PICS domain are preliminary in nature. Nevertheless, the summarised evidence paints

a promising picture of the development of eHealth interventions in this population. Future studies need to focus on larger-scale RCTs which will provide more insight into intervention effectiveness. The authors of the ICU-VR intervention have progressed to a larger RCT trial [59] in hope of generating more robust effects of the intervention on PICS outcomes. Other eHealth trials are also underway in this post-critical care phase of recovery [60–62]. Thus, whilst eHealth interventions can be concluded to be feasible, conclusions on effectiveness are premature at this point.

Even though no restriction was imposed on the language and country of article publication, the language used in the search strategy undoubtedly constrained its results. We acknowledge that if the search terms included other languages, other articles could be deemed eligible. This review adhered closely to the PICO framework [22] and search strings were systematically piloted in preliminary searches. The review attempted to be as broad as possible regarding the search strategy and the databases selected. Future research may also benefit from the inclusion of Medical Subject Headings (MeSH) terms to further expand the search.

## Conclusions

eHealth research and development in post-critical care rehabilitation is still early in its infancy with most studies focusing on feasibility. Based on the review findings, preliminary feasibility results are promising with research progressing to larger scale studies to derive more robust conclusions on effectiveness. Future research should be prioritised towards acceptability, targeting the cognitive domain, and exploring the effects of interventions targeting all 3 domains. eHealth is one vital solution in providing access, continuity, and sustainable care in the post-critical care setting.

## Abbreviations

ICU	Intensive Care Unit
PICS	Post Intensive Care Syndrome
HRQoL	Health Related Quality of Life
eHealth	Electronic Health
PRISMA	Preferred Reporting Items for Systematic Reviews and Meta-Analyses
PROSPERO	Prospective Register of Systematic Review databases
MMAT	Mixed Methods Appraisal Tool
RCT	Randomised Controlled Trial
TAU	Treatment as usual
Medical Subject Headings	MeSH

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13054-024-05089-6>.

**Additional file 1. Supplementary Material S1.** Database Search Strategies. **Supplementary Material S2.** Quality Assessment Criteria of the Mixed Method Appraisal Tool based on Study Design. **Table S3.** Feasibility outcomes and findings of eHealth interventions. **Table S4.** Acceptability Outcomes and findings of eHealth interventions

## Author contributions

All authors were involved with the conceptualisation of the review. DL wrote the draft and main manuscript. DL, ZL, and TD contributed to the article screening and inclusions. DL and ZL did data extraction independently with oversight from SS and TD. EJ, LD, SS, and TD provided critical feedback when reviewing and revising the manuscript. All authors reviewed the manuscript. All authors have read and approved the final manuscript.

## Funding

SS is funded by a MRC CARP award and a MRC Programme Grant MR/V037218/1 and by the NIHR/Wellcome Trust King's Clinical Research Facility and NIHR Maudsley Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. DL is funded by the NIHR Applied Research Collaboration Kent, Surrey and Sussex (NIHR ARC KSS). The views expressed are those of the author(s) and not necessarily those of the affiliated organisations.

## Availability of Data and Materials

Supplementary materials are provided and can be assessed online.

## Declarations

### Ethics Approval and Consent to Participate

Not applicable.

### Consent for Publication

Not applicable.

## Author details

<sup>1</sup>Canterbury Christ Church University, Canterbury, UK. <sup>2</sup>National Institute of Health and Care Research, Applied Research Collaboration Kent, Surrey and Sussex, UK. <sup>3</sup>School of Computing, University of Kent, Canterbury, UK. <sup>4</sup>College of Software and Big Data, Inner Mongolia Electronic Information Technical College, Hohhot, Inner Mongolia, China. <sup>5</sup>Maidstone and Tunbridge Wells NHS Trust, Kent, UK. <sup>6</sup>Kent and Medway Medical School, University of Kent and Canterbury Christ Church University, Canterbury, UK. <sup>7</sup>Centre for Health Services Studies, University of Kent, Canterbury, UK. <sup>8</sup>Institute of Psychiatry, Psychology, & Neuroscience, King's College London, London, UK.

Received: 24 June 2024 Accepted: 8 September 2024

Published online: 27 September 2024

## References

1. LaBuzetta JN, Rosand J, Vranceanu AM. post-intensive care syndrome: unique challenges in the neurointensive care unit. *Neurocrit Care*. 2019;31(3):534–45.
2. Zimmerman JE, Kramer AA, Knaus WA. Changes in hospital mortality for United States intensive care unit admissions from 1988 to 2012. *Crit Care*. 2013;17:1–9.
3. Prescott HC, Harrison DA, Rowan KM, Shankar-Hari M, Wunsch H. Temporal trends in mortality of critically ill patients with sepsis in the United Kingdom, 1988–2019. *Am J Respir Crit Care Med*. 2024;5:63.
4. Iwashyna TJ, Ely EW, Smith DM, Langa KM. Long-term cognitive impairment and functional disability among survivors of severe sepsis. *JAMA*. 2010;304(16):1787–94.
5. Needham DM, Davidson J, Cohen H, Hopkins RO, Weinert C, Wunsch H, Zawistowski C, Bemis-Dougherty A, Berney SC, Bienvenu OJ, Brady SL. Improving long-term outcomes after discharge from intensive care unit: report from a stakeholders' conference. *Crit Care Med*. 2012;40(2):502–9.
6. Bruno RR, Wernly B, Flaatten H, Fjølner J, Artigas A, Baldia PH, Binneboesel S, Bollen Pinto B, Schefold JC, Wolff G, Kelm M. The association of the Activities of Daily Living and the outcome of old intensive care patients suffering from COVID-19. *Ann Intensive Care*. 2022;12(1):26.
7. Gravante F, Trotta F, Latina S, Simeone S, Alvaro R, Vellone E, Pucciarelli G. Quality of life in ICU survivors and their relatives with post-intensive care syndrome: a systematic review. *Nurs Crit Care*. 2024;5:63.

8. Kamdar BB, Suri R, Suchyta MR, Digrande KF, Sherwood KD, Colantuoni E, Dinglas VD, Needham DM, Hopkins RO. Return to work after critical illness: a systematic review and meta-analysis. *Thorax*. 2020;75(1):17–27.
9. Brown SM, Bose S, Banner-Goodspeed V, Beesley SJ, Dinglas VD, Hopkins RO, Jackson JC, Mir-Kasimov M, Needham DM, Sevin CM. Approaches to addressing post-intensive care syndrome among intensive care unit survivors. A narrative review. *Ann Am Thor Soc*. 2019;16(8):947–56.
10. Nakanishi N, Liu K, Hatakeyama J, Kawauchi A, Yoshida M, Sumita H, Miyamoto K, Nakamura K. Post-intensive care syndrome follow-up system after hospital discharge: a narrative review. *J Intensive Care*. 2024;12(1):2.
11. Mehlhorn J, Freytag A, Schmidt K, Brunkhorst FM, Graf J, Troitzsch U, Schlattmann P, Wensing M, Gensichen J. Rehabilitation interventions for post-intensive care syndrome: a systematic review. *Crit Care Med*. 2014;42(5):1263–71.
12. Boehm LM, Danesh V, LaNoue M, Trochez RJ, Jones AC, Kimpel CC, Sevin CM. Factors influencing engagement with in-person intensive care unit recovery clinic services. *J Intensive Care Med*. 2023;38(4):375–81.
13. Black AD, Car J, Pagliari C, Anandan C, Cresswell K, Bokun T, Sheikh A. The impact of eHealth on the quality and safety of health care: a systematic overview. *PLoS Med*. 2011;8(1):e1000387.
14. Watanabe T, Ohsugi K, Suminaga Y, Somei M, Kikuyama K, Mori M, Maruo H, Kono N, Kotani T. An evaluation of the impact of the implementation of the Tele-ICU: a retrospective observational study. *J Intensive Care*. 2023;11(1):9.
15. Kalvelage C, Rademacher S, Dohmen S, Marx G, Benstoem C. Decision-making authority during tele-ICU care reduces mortality and length of stay—a systematic review and meta-analysis. *Crit Care Med*. 2021;49(7):1169–81.
16. Yuan C, Timmins F, Thompson DR. Post-intensive care syndrome: a concept analysis. *Int J Nurs Stud*. 2021;114(1):103814.
17. Morken IM, Storm M, Søreide JA, Urstad KH, Karlsen B, Husebø AM. Post-hospitalization follow-up of patients with heart failure using eHealth solutions: restricted systematic review. *J Med Internet Res*. 2022;24(2):e32946.
18. Hwang NK, Park JS, Chang MY. Telehealth interventions to support self-management in stroke survivors: a systematic review. *Healthcare*. 2021;9(4):52.
19. Zacharelou A, Major M, van der Meer P, van der Schaaf M, Vloet L, van Mol MM. Opinions and priorities for an e-health platform: a member consultation from an intensive care patient organisation. *Austr Crit Care*. 2024;2:59.
20. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, Chou R. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;29:372.
21. Lai D, Sukhi S, D'Oliveira T, Johnston E, Dikomitis L. A systematic review of the effectiveness of post-ICU eHealth interventions. PROSPERO 2023 CRD42023463036 Available from: [https://www.crd.york.ac.uk/prospero/display\\_record.php?ID=CRD42023463036](https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42023463036)
22. Schardt C, Adams MB, Owens T, Keitz S, Fontelo P. Utilization of the PICO framework to improve searching PubMed for clinical questions. *BMC Med Inform Decis Mak*. 2007;7:1–6.
23. Adams L, Valmaggia L, Simonoff E. Technology-based assessments and treatments of anxiety in autistic individuals: systematic review and narrative synthesis. *Rev J Aut Dev Disor*. 2022;9(4):571–95.
24. Hong QN, Fábregues S, Bartlett G, Boardman F, Cargo M, Dagenais P, Gagnon MP, Griffiths F, Nicolau B, O' Cathain A, Rousseau MC. The mixed methods appraisal tool (MMAT) version 2018 for information professionals and researchers. *Educ Inf*. 2018;34(4):285–91.
25. Pace R, Pluye P, Bartlett G, Macaulay AC, Salsberg J, Jagosh J, Seller R. Testing the reliability and efficiency of the pilot Mixed Methods Appraisal Tool (MMAT) for systematic mixed studies review. *Int J Nurs Stud*. 2012;49(1):47–53.
26. Balakrishnan B, Hamrick L, Alam A, Thompson J. Effects of COVID-19 acute respiratory distress syndrome intensive care unit survivor telemedicine clinic on patient readmission, pain perception, and self-assessed health scores: randomized, prospective, single-center, exploratory study. *JMIR Formative Research*. 2023;7(1):e43759.
27. Capin JJ, Jolley SE, Morrow M, Connors M, Hare K, MaWhinney S, Nordon-Craft A, Rauzi M, Flynn S, Stevens-Lapsley JE, Erlandson KM. Safety, feasibility and initial efficacy of an app-facilitated telerehabilitation (AFTER) programme for COVID-19 survivors: a pilot randomised study. *BMJ Open*. 2022;12(7):e061285.
28. Cox CE, Hough CL, Jones DM, Ungar A, Reagan W, Key MD, Gremore T, Olsen MK, Sanders L, Greeson JM, Porter LS. Effects of mindfulness training programmes delivered by a self-directed mobile app and by telephone compared with an education programme for survivors of critical illness: a pilot randomised clinical trial. *Thorax*. 2019;74(1):33–42.
29. Denehy L, Berney S, Whitburn L, Edbrooke L. Quantifying physical activity levels of survivors of intensive care: a prospective observational study. *Phys Ther*. 2012;92(12):1507–17.
30. Estrup S, Kjer CK, Vilhelmsen F, Poulsen LM, Gøgenur I, Mathiesen O. Physical function and actigraphy in intensive care survivors—a prospective 3-month follow-up cohort study. *Acta Anaesthesiol Scand*. 2019;63(5):647–52.
31. Jackson JC, Ely EW, Morey MC, Anderson VM, Denne LB, Clune J, Siebert CS, Archer KR, Torres R, Janz D, Schiro E. Cognitive and physical rehabilitation of intensive care unit survivors: results of the RETURN randomized controlled pilot investigation. *Crit Care Med*. 2012;40(4):1088–97.
32. Kovaleva MA, Jones AC, Kimpel CC, Lauderdale J, Sevin CM, Stollings JL, Jackson JC, Boehm LM. Patient and caregiver experiences with a telemedicine intensive care unit recovery clinic. *Heart Lung*. 2023;1(58):47–53.
33. Park N, Oberlin L, Cherestal S, Castellano CB, Dargis M, Wyka KE, Jaywant A, Kanellopoulos D. Trajectory and outcomes of psychiatric symptoms in first-wave COVID-19 survivors referred for telepsychotherapy. *Gen Hosp Psychiatry*. 2023;81:86.
34. Parker AM, Nelliot A, Chessare CM, Malik AM, Koneru M, Hosey MM, Ozok AA, Lyons KD, Needham DM. Usability and acceptability of a mobile application prototype for a combined behavioural activation and physical rehabilitation intervention in acute respiratory failure survivors. *Aust Crit Care*. 2020;33(6):511–7.
35. Rose L, Apps C, Brooks K, Terblanche E, Hart N, Meyer J. Novel digitally enabled care pathway to support postintensive care recovery and goal attainment following critical illness. *BMJ Innov*. 2022;8(1):63.
36. Vlaker JH, Van Bommel J, Wils EJ, Korevaar TI, Bienvenu OJ, Klijn E, Gommers D, Van Genderen ME. Virtual reality to improve sequelae of the post-intensive care syndrome: a multicenter, randomized controlled feasibility study. *Crit Care Explor*. 2021;3(9):e0538.
37. Vlaker JH, van Bommel J, Wils EJ, Bienvenu J, Hellemons ME, Korevaar TI, Schut AF, Labout JA, Schreuder LL, van Bavel MP, Gommers D. Intensive care unit-specific virtual reality for critically ill patients with COVID-19: multicenter randomized controlled trial. *J Med Internet Res*. 2022;24(1):e32368.
38. Wood MD, Maslove DM, Muscedere J, Scott SH, Boyd JG, Canadian Critical Care Trials Group. Robotic technology provides objective and quantifiable metrics of neurocognitive functioning in survivors of critical illness: a feasibility study. *J Crit Care*. 2018;48:228–36.
39. Major ME, Kwakman R, Kho ME, Connolly B, McWilliams D, Denehy L, Hanekom S, Patman S, Gosselin K, Jones C, Nollet F. Surviving critical illness: what is next? An expert consensus statement on physical rehabilitation after hospital discharge. *Crit Care*. 2016;20:1.
40. Spies CD, Krampe H, Paul N, Denke C, Kiselev J, Piper SK, Kruppa J, Grunow JJ, Steinecke K, Gülmez T, Scholtz K. Instruments to measure outcomes of post-intensive care syndrome in outpatient care settings—results of an expert consensus and feasibility field test. *J Intens Care Soc*. 2021;22(2):159–74.
41. Needham DM, Sepulveda KA, Dinglas VD, Chessare CM, Friedman LA, Bingham CO III, Turnbull AE. Core outcome measures for clinical research in acute respiratory failure survivors. An international modified Delphi consensus study. *Am J Respir Crit Care Med*. 2017;196(9):1122–30.
42. Dinglas VD, Cherukuri SP, Needham DM. Core outcomes sets for studies evaluating critical illness and patient recovery. *Curr Opin Crit Care*. 2020;26(5):489–99.
43. Connolly B, Salisbury L, O'Neill B, Geneen LJ, Douiri A, Grocott MP, ERACIP Group. Exercise rehabilitation following intensive care unit discharge for recovery from critical illness. *Cochr Database Syst Rev*. 2015;6:35.
44. Kress JP. Sedation and mobility: changing the paradigm. *Crit Care Clin*. 2013;29(1):67–75.
45. Baker LD, Frank LL, Foster-Schubert K, Green PS, Wilkinson CW, McTiernan A, Plymate SR, Fishel MA, Watson GS, Cholerton BA, Duncan GE. Effects

- of aerobic exercise on mild cognitive impairment: a controlled trial. *Arch Neurol.* 2010;67(1):71–9.
46. Heesakkers H, Van Der Hoeven JG, Corsten S, Janssen I, Ewalds E, Simons KS, Westerhof B, Rettig TC, Jacobs C, Van Santen S, Slooter AJ. Clinical outcomes among patients with 1-year survival following intensive care unit treatment for COVID-19. *JAMA.* 2022;327(6):559–65.
  47. Marra A, Pandharipande PP, Girard TD, Patel MB, Hughes CG, Jackson JC, Thompson JL, Chandrasekhar R, Ely EW, Brummel NE. Co-occurrence of post-intensive care syndrome problems among 406 survivors of critical illness. *Crit Care Med.* 2018;46(9):1393–401.
  48. Kang J, Jeong YJ, Hong J. The effect of postintensive care syndrome on the quality of life of intensive care unit survivors: a secondary analysis. *Aust Crit Care.* 2021;34(3):246–53.
  49. Skivington K, Matthews L, Simpson SA, Craig P, Baird J, Blazeby JM, Boyd KA, Craig N, French DP, McIntosh E, Petticrew M. A new framework for developing and evaluating complex interventions: update of Medical Research Council guidance. *BMJ.* 2021;6:374.
  50. Prestwich A, Sniehotta FF, Whittington C, Dombrowski SU, Rogers L, Michie S. Does theory influence the effectiveness of health behavior interventions? Meta-analysis. *Health Psychol.* 2014;33(5):465.
  51. Hawkins J, Madden K, Fletcher A, Midgley L, Grant A, Cox G, Moore L, Campbell R, Murphy S, Bonell C, White J. Development of a framework for the co-production and prototyping of public health interventions. *BMC Public Health.* 2017;17:1–1.
  52. Greenhalgh T, Shaw S, Wherton J, Vijayaraghavan S, Morris J, Bhattacharya S, Hanson P, Campbell-Richards D, Ramoutar S, Collard A, Hodkinson I. Real-world implementation of video outpatient consultations at macro, meso, and micro levels: mixed-method study. *J Med Internet Res.* 2018;20(4):e150.
  53. Varsi C, Solberg Nes L, Kristjansdottir OB, Kelders SM, Stenberg U, Zangi HA, Børusund E, Weiss KE, Stubhaug A, Asbjørnsen RA, Westeng M. Implementation strategies to enhance the implementation of eHealth programs for patients with chronic illnesses: realist systematic review. *J Med Internet Res.* 2019;21(9):e14255.
  54. Waltz TJ, Powell BJ, Matthieu MM, Damschroder LJ, Chinman MJ, Smith JL, Proctor EK, Kirchner JE. Use of concept mapping to characterize relationships among implementation strategies and assess their feasibility and importance: results from the Expert Recommendations for Implementing Change (ERIC) study. *Implement Sci.* 2015;10:1–8.
  55. Zaman SB, Khan RK, Evans RG, Thrift AG, Maddison R, Islam SM. Exploring barriers to and enablers of the adoption of information and communication technology for the care of older adults with chronic diseases: scoping review. *JMIR Agi.* 2022;5(1):e25251.
  56. Hasnan S, Aggarwal S, Mohammadi L, Koczwara B. Barriers and enablers of uptake and adherence to digital health interventions in older patients with cancer: a systematic review. *J Geriatr Oncol.* 2022;13(8):1084–91.
  57. Kampmeijer R, Pavlova M, Tambor M, Golinowska S, Groot W. The use of e-health and m-health tools in health promotion and primary prevention among older adults: a systematic literature review. *BMC Health Serv Res.* 2016;16:467–79.
  58. Wilson J, Heinsch M, Betts D, Booth D, Kay-Lambkin F. Barriers and facilitators to the use of e-health by older adults: a scoping review. *BMC Public Health.* 2021;21:1–2.
  59. Vlaker JH, van Bommel J, Wils EJ, Korevaar TI, Taccone F, Schut AF, Elderman JH, Labout JA, Raben AM, Dijkstra A, Achterberg S. Effect of intensive care unit-specific virtual reality (ICU-VR) to improve psychological well-being in ICU survivors: study protocol for an international, multi-centre, randomised controlled trial—the HORIZON-IC study. *BMJ Open.* 2022;12(9):e061876.
  60. Turan Z, Topaloglu M, Ozyemisci TO. Is tele-rehabilitation superior to home exercise program in COVID-19 survivors following discharge from intensive care unit?—a study protocol of a randomized controlled trial. *Physiother Res Int.* 2021;26(4):e1920.
  61. Limotai C, Ingsathit A, Thadanipon K, Pattanapruteep O, Pattanateepaporn A, Phanthumchinda K, Suwanwela NC, Thaipisuttikul I, Boonyapisit K, Thakkinstian A. Efficacy and economic evaluation of delivery of care with tele-continuous EEG in critically ill patients: a multicentre, randomised controlled trial (Tele-cRCT) study protocol. *BMJ Open.* 2020;10(3):e033195.
  62. Ewens B, Myers H, Whitehead L, Seaman K, Sundin D, Hendricks J. A web-based recovery program (ICUTogether) for intensive care survivors: protocol for a randomized controlled trial. *JMIR Res Prot.* 2019;8(1):e10935.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.