

COMMENTARY

Are arterial blood gases necessary in the evaluation of acutely dyspneic patients?

Christopher Junker^{1*} and Guillermo Gutierrez²

See related research by Burri *et al.*, <http://ccforum.com/content/15/3/R145>

Abstract

Arterial blood gases (ABG) are obtained commonly in dyspneic persons presenting to emergency departments. The study by Burri and colleagues found that the information contained in ABG fails to distinguish between pulmonary and other causes of dyspnea. On the other hand, arterial pH was highly predictive of ICU admission and outcome. Until large clinical studies show equivalence between peripheral venous and ABG, we will continue to advocate the use of ABG in the evaluation of acute dyspnea.

The recent study by Burri and colleagues examined the usefulness of arterial blood gases (ABG) in the diagnosis and prognosis of 530 dyspneic patients with ABG drawn upon presentation to the emergency department [1]. The study was a retrospective analysis of prospectively collected data performed at a single center. They concluded that ABG cannot be used to distinguish between pulmonary and other causes of dyspnea.

Prior to the introduction of automated blood gas analyzers, blood PO₂ and PCO₂ could be obtained only by laborious and often inaccurate laboratory methods, such as gas tonometry and Van Slyke manometric extraction of plasma total carbon dioxide. Improvements in electrode technology came with the development of the Astrup pH electrode [2], the Stow/Severinghaus-type PCO₂ electrode [3,4] and the polarographic oxygen Clark electrode [5]. Although pulse oximetry has largely superseded the use of arterial PO₂ (PaO₂), the ease by which ABG can now be measured has made this test *de rigueur* in the workup of dyspneic patients.

We agree with Burri and colleagues in that PaO₂ values serve mainly to triage and to guide treatment, rather than to differentiate among the causes of dyspnea. Except for anxiety hyperventilation, it is difficult to rely solely on ABG to identify a specific clinical syndrome. This is not new information. The poor predictive value of PaO₂ in diagnosing patients with pulmonary embolism is well known. In the PIOPED I study of patients with angiographically proven pulmonary embolism and no prior cardiopulmonary disease, 26% had PaO₂ >80 mmHg [6]. This figure was 38% in the PIOPED II study (*n* = 42) [7,8]. The probability of diagnosing an acute pulmonary embolism based on changes in PaO₂ did not achieve statistical significance. Further, no combination of PO₂ and PCO₂ values could reliably exclude pulmonary embolism.

Burri and colleagues also report that arterial pH was a significant predictor of short-term and long-term outcome. Multiple physiological buffers act to preserve the concentration of hydrogen ions in blood within a relatively small range. Decreases in arterial pH are likely to reflect severe impairments or even exhaustion of systemic compensatory mechanisms in patients with acute dyspnea. The predictive value of the arterial pH in dyspneic patients noted by Burri and colleagues supports this hypothesis.

Burri and colleagues propose the use of venous blood pH in the initial evaluation of acute dyspnea, based on data from several single-center studies reporting relatively close limits of agreement (−0.11 to +0.04) between arterial pH and venous blood pH [9,10]. One must keep in mind, however, that muscular activity or regional microcirculatory alterations could result in misleadingly low regional pH values in septic or agitated patients.

Substituting venous blood pH for arterial pH in the evaluation of dyspneic patients is an appealing notion, but one that requires testing in multicenter, prospective clinical studies. Until these studies corroborate the equivalence between arterial and peripheral venous blood gases, or until non-invasive techniques to monitor arterial PCO₂ and pH become available, we shall continue to support the use of ABG in the initial evaluation and treatment of patients with acute dyspnea.

*Correspondence: cjunker@mfa.gwu.edu

¹Department of Anesthesiology and Critical Care Medicine, The George Washington University MFA, 2150 Pennsylvania Ave, NW, Washington, DC 20037, USA

Full list of author information is available at the end of the article

Abbreviations

ABG, arterial blood gases; ICU, intensive care unit; PaCO₂, arterial partial pressure of carbon dioxide; PCO₂, partial pressure of carbon dioxide; PO₂, partial pressure of oxygen.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Anesthesiology and Critical Care Medicine, The George Washington University MFA, 2150 Pennsylvania Ave, NW, Washington, DC 20037, USA. ²Department of Medicine and Anesthesiology, Division of Pulmonary and Critical Care Medicine, The George Washington University MFA, 2150 Pennsylvania Ave, NW, Washington, DC 20037, USA.

Published: 2 August 2011

References

1. Burri E, Potocki M, Drexler B, Schuetz P, Mebazaa A, Ahlfeld U, Balmelli C, Heinisch C, Noveanu M, Breidhardt T, Schaub N, Reichlin T, Mueller C: **Value of arterial blood gas analysis in patients with acute dyspnea: an observational study.** *Crit Care* 2011, **15**:R145.
2. Astrup P, Schröder S: **Apparatus for anaerobic determination of pH in blood.** *Scand J Clin Invest* 1956, **8**:30-32.
3. Stow RW, Baer RF, Randall B: **Rapid measurement of the tension of carbon dioxide in blood.** *Arch Phys Med Rehabil* 1957, **38**:646-650.
4. Severinghaus JW, Bradley AF: **Electrodes for blood PO₂ and PCO₂ determination.** *J Appl Physiol* 1958, **13**:515-520.
5. Clark LC Jr: **Monitor and control of blood and tissue oxygen tensions.** *Trans Am Soc Artif Intern Organs* 1956, **2**:41-48.
6. PIOPED Investigators: **Value of the ventilation/perfusion scan in acute pulmonary embolism. Results of the prospective investigation of pulmonary embolism diagnosis (PIOPED).** *JAMA* 1990, **263**:2753-2755.
7. Stein PD, Fowler SE, Goodman LR, Gottschalk A, Hales CA, Hull RD, LEEPER KV Jr, Popovich J Jr, Quinn DA, Sos TA, Sostman HD, Tapson VF, Wakefield TW, Weg JG, Woodard PK; PIOPED II Investigators: **Multidetector computed tomography for acute pulmonary embolism.** *N Engl J Med* 2006, **354**:2317-2327.
8. Stein PD, Beemath A, Matta F, Weg JG, Yusen RD, Hales CA, Hull RD, LEEPER KV Jr, Sostman HD, Tapson VF, Buckley JD, Gottschalk A, Goodman LR, Wakefield TW, Woodard PK: **Clinical characteristics of patients with acute pulmonary embolism: data from PIOPED II.** *Am J Med* 2007, **120**:871-879.
9. Kelly AM, McAlpine R, Kyle E: **Venous pH can safely replace arterial pH in the initial evaluation of patients in the emergency department.** *Emerg Med J* 2001, **18**:340-342.
10. Treger R, Pirouz S, Kamangar N, Corry D: **Agreement between central venous and arterial blood gas measurements in the intensive care unit.** *Clin J Am Soc Nephrol* 2010, **5**:390-394.

doi:10.1186/cc10279

Cite this article as: Junker C, Gutierrez G: **Are arterial blood gases necessary in the evaluation of acutely dyspneic patients?** *Critical Care* 2011, **15**:176.