

LETTER

Preventing deaths related to percutaneous tracheostomy: safety is never too much!

Luca Cabrini*, Massimiliano Greco, Laura Pasin, Giacomo Monti, Sergio Colombo and Alberto Zangrillo

See related research by Simon et al., <http://ccforum.com/content/17/5/R258>

We read with great interest the study by Simon and colleagues about intra- and post-procedural mortality related to percutaneous dilatational tracheostomy (PDT) [1]. We appreciated the huge efforts made by the authors to collect valuable data on this rare event: their findings can significantly improve daily practice in PDT performance and management in the ICU. We present three additional comments.

First of all, the authors affirmed that a standard use of ultrasound neck evaluation and continuous bronchoscopy could reduce the incidence of bleeding and airway complications, respectively. However, it must be underlined that the most common intra-procedural cause of death reported in this study was pneumothorax. Thus, the complete or almost complete tracheal occlusion by dilators or the bronchoscope should be minimized to avoid air trapping. Moreover, the safest ventilatory setting while PDT is performed is unknown, and research on this topic is urgently needed. Air trapping within the lungs (a potential cause of overinflation and pneumothorax), caused by a valve effect while performing PDT, should be carefully avoided through the adoption of lower positive end-expiratory pressure, lower respiratory rate, and smaller tidal volumes [2].

Secondly, the dislocation of the tracheal cannula is a common cause of late mortality: the development of a dedicated 'crisis' flowchart, the immediate availability of the required equipment, and periodic personnel retraining drills [3] should be considered similarly to other protocols already validated for emergent conditions like intubation in critically ill patients [4].

Finally, the available PDT techniques are not equally safe. Some evidence exists that the single-dilator technique is safer than the others at least intra-procedurally and could be considered the first choice [5].

Abbreviation

PDT: percutaneous tracheostomy.

Competing interests

The authors declare that they have no competing interests.

Published: 13 Feb 2014

References

1. Simon M, Metschke M, Braune SA, Püschel K, Kluge S: **Death after percutaneous dilatational tracheostomy: a systematic review and analysis of risk factors.** *Crit Care* 2013, **17**:R258.
2. Sheu CC, Tsai JR, Cheng MH, Chong IW, Huang MS, Hwang JJ: **Safety of performing percutaneous dilatational tracheostomy in patients with preexisting barotrauma.** *Kaohsiung J Med Sci* 2006, **22**:570-574.
3. Thomas AN, McGrath BA: **Patient safety incidents associated with airway devices in critical care: reports to the UK National Patient Safety Agency.** *Anaesthesia* 2009, **64**:358-365.
4. Jaber S, Jung B, Corne P, Sebbane M, Muller L, Chanques G, Verzilli D, Jonquet O, Eledjam JJ, Lefrant JY: **An intervention to decrease complications related to endotracheal intubation in the intensive care unit: a prospective, multiple-center study.** *Intensive Care Med* 2010, **36**:248-255.
5. Cabrini L, Monti G, Landoni G, Biondi-Zoccai G, Boroli F, Mamo D, Plumari VP, Colombo S, Zangrillo A: **Percutaneous tracheostomy, a systematic review.** *Acta Anaesthesiol Scand* 2012, **56**:270-281.

10.1186/cc13727

Cite this article as: Cabrini et al.: Preventing deaths related to percutaneous tracheostomy: safety is never too much!. *Critical Care* 2014, **18**:406

* Correspondence: cabrini.luca@hsr.it
Department of Anesthesia and Intensive Care, San Raffaele Scientific Institute,
Via Olgettina 60, 20132 Milan, Italy