## Correction

## Correction: Bispectral index versus COMFORT score to determine the level of sedation in paediatric intensive care unit patients: a prospective study

Andreas E Triltsch<sup>1</sup>, Grit Nestmann<sup>2</sup>, Helmut Orawa<sup>3</sup>, Maryam Moshirzadeh<sup>4</sup>, Michael Sander<sup>4</sup>, Joachim Große<sup>4</sup>, Arka Genähr<sup>4</sup>, Wolfgang Konertz<sup>5</sup>, Claudia D Spies<sup>4</sup>

Corresponding author: Claudia Spies, claudia.spies@charite.de

Published online: 2 September 2005

This article is online at http://ccforum.com/content/9/5/426

© 2005 Spies et al., licensee BioMed Central Ltd.

Critical Care 2005, 9:426 (DOI 10.1186/cc3810)

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/2.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is cited.

After the publication of this article [1] we noticed that the affiliation details were incorrect and should be as follows:

Andreas E Triltsch – Department of Anesthesiology and Intensive Care Medicine, Campus Benjamin Franklin, Charité University Hospital Berlin, Berlin, Germany

Grit Nestmann - Department of Pediatrics, Campus Benjamin Franklin, Charité University Hospital Berlin, Berlin, Germany

Helmut Orawa – Department of Medical Informatics, Biometry, and Epidemiology, Charité University Medicine Berlin, Berlin, Germany

Maryam Moshirzadeh, Michael Sander, Joachim Große and Claudia D Spies - Department of Anesthesiology and Intensive Care Medicine, Campus Charité Mitte, Charité University Hospital Berlin, Berlin, Germany

Wolfgang Konertz - Department of Cardiac Surgery, Campus Charité Mitte, Charité University Hospital Berlin, Berlin, Germany

## References

 Triltsch AE, Nestmann G, Orawa H, Moshirzadeh M, Sander M, Große J, Genähr A, Konertz W, Spies CD: Bispectral index versus COMFORT score to determine the level of sedation in paediatric intensive care unit patients: a prospective study. Crit Care 2005, 9:R9-R17

<sup>&</sup>lt;sup>1</sup>Department of Anesthesiology and Intensive Care Medicine, Campus Benjamin Franklin, Charité University Hospital Berlin, Berlin, Germany

<sup>&</sup>lt;sup>2</sup>Department of Pediatrics, Campus Benjamin Franklin, Charité University Hospital Berlin, Berlin, Germany

<sup>&</sup>lt;sup>3</sup>Department of Medical Informatics, Biometry and Epidemiology, Charité University Medicine Berlin, Berlin, Germany

<sup>&</sup>lt;sup>4</sup>Department of Anesthesiology and Intensive Care Medicine, Campus Charité Mitte, Charité University Hospital Berlin, Berlin, Germany

<sup>&</sup>lt;sup>5</sup>Department of Cardiac Surgery, Campus Charité Mitte, Charité University Hospital Berlin, Berlin, Germany