

## Correction

# Correction: Management of bleeding following major trauma: a European guideline

Donat R Spahn<sup>1</sup>, Vladimir Cerny<sup>2</sup>, Timothy J Coats<sup>3</sup>, Jacques Duranteau<sup>4</sup>, Enrique Fernández-Mondéjar<sup>5</sup>, Giovanni Gordini<sup>6</sup>, Philip F. Stahel<sup>7</sup>, Beverley J Hunt<sup>8</sup>, Radko Komadina<sup>9</sup>, Edmund Neugebauer<sup>10</sup>, Yves Ozier<sup>11</sup>, Louis Riddez<sup>12</sup>, Arthur Schultz<sup>13</sup>, Jean-Louis Vincent<sup>14</sup>, Rolf Rossaint<sup>15</sup>

<sup>1</sup>Department of Anesthesiology, University Hospital Zurich, Rämistrasse 100, 8091 Zurich, Switzerland

<sup>2</sup>Charles University in Prague, Faculty of Medicine in Hradec Králové, Department of Anaesthesiology and Intensive Care Medicine, University Hospital Hradec Králové, Sokolska 581, 50005 Hradec Králové, Czech Republic

<sup>3</sup>Leicester Royal Infirmary, Accident and Emergency Department, Infirmary Square, Leicester LE1 5WW, UK

<sup>4</sup>Department of Anaesthesia and Intensive Care, University of Paris XI Faculté de Médecine Paris-Sud, 63 rue Gabriel Péri, 94276 Le Kremlin-Bicêtre, France

<sup>5</sup>Department of Emergency and Critical Care Medicine, University Hospital Virgen de las Nieves, ctra de Jaén s/n, 18013 Granada, Spain

<sup>6</sup>Department of Anaesthesia and Intensive Care, Ospedale Maggiore, Largo Nigrisoli 2, 40100 Bologna, Italy

<sup>7</sup>Department of Orthopaedic Surgery, Denver Health Medical Center, University of Colorado Medical School, 777 Bannock Street, Denver, CO 80204, USA

<sup>8</sup>Departments of Haematology, Pathology and Rheumatology, Guy's & St Thomas' Foundation Trust, Lambeth Palace Road, London SE1 7EH, UK

<sup>9</sup>Department of Traumatology, General and Teaching Hospital Celje, 3000 Celje, Slovenia

<sup>10</sup>Institute for Research in Operative Medicine, University of Witten/Herdecke, Ostmerheimerstrasse 200, 51109 Köln (Merheim), Germany

<sup>11</sup>Department of Anaesthesia and Intensive Care, Université René Descartes Paris 5, AP-HP, Hôpital Cochin, 27 rue du Fbg Saint-Jacques, 75014 Paris, France

<sup>12</sup>Department of Surgery and Trauma, Karolinska University Hospital, 171 76 Solna, Sweden

<sup>13</sup>Ludwig-Boltzmann-Institute for Experimental and Clinical Traumatology, Donaueschingenstrasse 13, 1200 Vienna, Austria

<sup>14</sup>Department of Intensive Care, Erasme Hospital, University of Brussels, Belgium, route de Lennik 808, 1070 Brussels, Belgium

<sup>15</sup>Department of Anaesthesiology, University Hospital Aachen, Pauwelsstraße 30, 52074 Aachen, Germany

Corresponding author: Rolf Rossaint, [rossaint@post.rwth-aachen.de](mailto:rossaint@post.rwth-aachen.de)

Published: 24 April 2007

*Critical Care* 2007, **11**:414 (doi:10.1186/cc5781)

This article is online at <http://ccforum.com/content/11/2/414>

© 2007 BioMed Central Ltd

It has been brought to our attention that the seventh sentence from paragraph two of Recommendation 17 in our research article [1], published in February 2007, is incorrect.

"A clinical trial with brain injury patients found that hypertonic saline reduces intracranial pressure more effectively than dextran solution with 20% mannitol"

This should read:

"A clinical trial with brain injury patients found that hypertonic saline with dextran reduces intracranial pressure more effectively than 20% mannitol solution when compared in equimolar dosing"

## Reference

1. Spahn DR, Cerny V, Coats TJ, Fernandez-Mondejar, Duranteau J, Hunt BJ, Komadina R, Neugebauer E, Ozier Y, Riddez L, Schultz A, Vincent JL, Rossaint R: **Management of bleeding following major trauma: a European guideline.** *Critical Care* 2007, **11**:R17.