RESEARCH

Open Access

Role of selective V₂-receptor-antagonism in septic shock: a randomized, controlled, experimental study

Sebastian Rehberg^{1*}, Christian Ertmer¹, Matthias Lange¹, Andrea Morelli², Elbert Whorton³, Martin Dünser⁴, Anne-Katrin Strohacker¹, Erik Lipke¹, Tim G Kampmeier¹, Hugo Van Aken¹, Daniel L Traber⁵, Martin Westphal¹

Abstract

Introduction: V₂-receptor (V₂R) stimulation potentially aggravates sepsis-induced vasodilation, fluid accumulation and microvascular thrombosis. Therefore, the present study was performed to determine the effects of a first-line therapy with the selective V₂R-antagonist (Propionyl₁-D-Tyr(Et)₂-Val₄-Abu₆-Arg_{8,9})-Vasopressin on cardiopulmonary hemodynamics and organ function vs. the mixed V_{1a}R/V₂R-agonist arginine vasopressin (AVP) or placebo in an established ovine model of septic shock.

Methods: After the onset of septic shock, chronically instrumented sheep were randomly assigned to receive first-line treatment with the selective V₂R-antagonist (1 µg/kg per hour), AVP (0.05 µg/kg per hour), or normal saline (placebo, each *n* = 7). In all groups, open-label norepinephrine was additionally titrated up to 1 µg/kg per minute to maintain mean arterial pressure at 70 ± 5 mmHg, if necessary.

Results: Compared to AVP- and placebo-treated animals, the selective V₂R-antagonist stabilized cardiopulmonary hemodynamics (mean arterial and pulmonary artery pressure, cardiac index) as effectively and increased intravascular volume as suggested by higher cardiac filling pressures. Furthermore, left ventricular stroke work index was higher in the V₂R-antagonist group than in the AVP group. Notably, metabolic (pH, base excess, lactate concentrations), liver (transaminases, bilirubin) and renal (creatinine and blood urea nitrogen plasma levels, urinary output, creatinine clearance) dysfunctions were attenuated by the V₂R-antagonist when compared with AVP and placebo. The onset of septic shock was associated with an increase in AVP plasma levels as compared to baseline in all groups. Whereas AVP plasma levels remained constant in the placebo group, infusion of AVP increased AVP plasma levels up to 149 ± 21 pg/mL. Notably, treatment with the selective V₂R-antagonist led to a significant decrease of AVP plasma levels as compared to shock time (*P* < 0.001) and to both other groups (*P* < 0.05 vs. placebo; *P* < 0.001 vs. AVP). Immunohistochemical analyses of lung tissue revealed higher hemeoxygenase-1 (vs. placebo) and lower 3-nitrotyrosine concentrations (vs. AVP) in the V₂R-antagonist group. In addition, the selective V₂R-antagonist slightly prolonged survival (14 ± 1 hour) when compared to AVP (11 ± 1 hour, *P* = 0.007) and placebo (11 ± 1 hour, *P* = 0.025).

Conclusions: Selective V₂R-antagonism may represent an innovative therapeutic approach to attenuate multiple organ dysfunction in early septic shock.

Introduction

Arginine vasopressin (AVP) is recommended by the Surviving Sepsis Campaign to 'be subsequently added to norepinephrine' in volume- and catecholamine-refractory septic shock [1]. In the randomized, controlled, multicenter Vasopressin and Septic Shock Trial (VASST), however,

AVP failed to reduce overall mortality as compared with norepinephrine among patients with septic shock [2].

AVP represents a mixed V_{1a}/V₂ receptor (V_{1a}R/V₂R) agonist with a selectivity of 1:1 for each of these receptors. Whereas particular attention has been paid to the vasoconstriction mediated by vascular V_{1a}Rs [3,4], there is increasing evidence that stimulation of extrarenal (endothelial) V₂Rs [5-7] may aggravate sepsis-induced vasodilation [4,8], fluid accumulation [9], leukocyte rolling [10], and microvascular thrombosis [11]. Against this

* Correspondence: Sebastian_Rehberg@web.de

¹Department of Anesthesiology and Intensive Care, University of Muenster, Albert-Schweitzer-Str. 33, Muenster 48149, Germany

Full list of author information is available at the end of the article

background, selective V₂R-antagonism potentially represents a new therapeutic approach in septic shock.

We hypothesized that a first-line therapy with the selective V₂R-antagonist (propionyl₁-D-Tyr(Et)₂-Val₄-Abu₆-Arg_{8,9}) vasopressin [12,13] is more effective than infusion of placebo and AVP in restoring cardiovascular and renal functions in early ovine septic shock. Open-label norepinephrine was additionally titrated to maintain mean arterial pressure (MAP) in each group if necessary. Therefore, the present study was designed as a prospective, randomized, controlled, laboratory experiment to elucidate the effects of these treatment strategies on cardiopulmonary hemodynamics, mesenteric blood flow, global oxygen transport, acid-base balance, organ function, AVP plasma levels, oxidative stress, and mortality. The study hypothesis was tested in an established ovine model of fulminant septic shock resulting from generalized fecal peritonitis [14,15].

Materials and methods

Instrumentation and surgical procedures

After approval by the Local Animal Research Committee, 21 female sheep were anesthetized, mechanically ventilated, and instrumented for chronic hemodynamic monitoring using an established protocol [14,15]. Details on the instrumentation and surgical procedures are provided in the supplemental digital content in Additional file 1.

Experimental protocol

Following baseline (BL) measurements, autologous feces were injected into the peritoneal cavity via an intraperitoneal suction catheter. When septic shock had developed (so-called 'shock time' [ST], defined as MAP of less than 60 mm Hg), a second set of measurements was performed. The animals were then randomly assigned to receive a first-line therapy with the selective V₂R-antagonist (1 µg/kg per hour; *n* = 7; Bachem Distribution Services AG, Weil am Rhein, Germany), AVP (0.05 µg/kg per hour, equivalent to 0.5 mU/kg per minute or 0.035 U/minute in a 70-kg patient; *n* = 7; American Regent Inc., Shirley, NY, USA), or normal saline (*n* = 7; B. Braun Melsungen AG, Melsungen, Germany). Open-label norepinephrine (Arterenol; Aventis Pharma, Frankfurt, Germany) was additionally titrated up to 1 µg/kg per minute to maintain MAP at 70 ± 5 mm Hg in all groups, if necessary. To ensure normovolemia, continuous infusions of balanced isotonic crystalloids (Sterofundin ISO; B. Braun Melsungen AG, Melsungen, Germany) and 6% hydroxyethyl starch 130/0.4 (Voluven; Fresenius Kabi, Bad Homburg, Germany) were infused at 8 and 4 mL/kg per hour, respectively, after ST. Additional fluids (crystalloid/colloid ratio of 2:1) were infused if

hematocrit exceeded BL values during the 24-hour study period [14].

Hemodynamic measurement, blood gas, laboratory, and histological analyses

Hemodynamic measurements, arterial and mixed venous blood gas, and laboratory analyses of variables of organ dysfunction and AVP plasma levels were performed at specific time points. Details on these measurements are provided in the supplemental digital content in Additional file 1.

Immunohistochemical analyses

Following death, tissue samples were immediately stored for immunohistochemical analyses. Pulmonary concentrations of hemeoxygenase-1 (StressXpress Human HO-1 ELISA [enzyme-linked immunosorbent assay] Kit; Stressgen Bioreagents, Ann Arbor, MI, USA) and 3-nitrotyrosine (Hycult biotechnology 3-nitrotyrosine solid-phase ELISA; Cell Sciences, Canton, MA, USA) were determined as described previously [16,17].

Statistical analyses

Sigma Stat 3.1 software (Systat Software, Inc., San Jose, CA, USA) was used for statistical analyses. Analysis-of-variance methodologies appropriate for two-factor experiments with repeated measures across time for each animal were used. Each variable was analyzed separately for differences among groups and differences across time and for group by time interaction. After confirmation of the significance of different group effects over time, *post hoc* pairwise comparisons among groups were performed using the Student-Newman-Keuls procedure to adjust for the elevated false-positive rate found otherwise in multiple testing. After 10 hours, no statistical analyses were performed, because the small number of animals alive in the placebo and the AVP group did not allow reliable testing anymore. Survival times were calculated using a log-rank test. Group differences were analyzed by pairwise multiple comparison with the Holm-Sidak test. Differences were considered statistically significant for *P* values of less than 0.05.

Results

Baseline characteristics

There were no differences among study groups in any of the investigated variables at BL and ST. Mean body weight (37 ± 1 kg) and time to onset of septic shock (7 ± 1 hours) did not differ between groups.

Cardiopulmonary hemodynamics

Changes in cardiopulmonary variables are presented in Figures 1 and 2 and Table 1. Septic shock was characterized by decreases in MAP, systemic vascular resistance

index, and left ventricular stroke work index (LVSWI) (ST: $P < 0.001$ versus BL each). All three treatment strategies maintained MAP within the target range of 70 ± 5 mm Hg for the first 4 hours after ST (4 hours: $P < 0.01$ versus ST each; Table 1). However, after the dose limitation for norepinephrine had been reached, MAP and systemic vascular resistance index fell significantly below ST values in all groups (10 hours: $P < 0.05$ versus ST each; Table 1). There were no statistically significant differences in cumulative norepinephrine requirements among study groups (Figure 1a).

LVSWI increased significantly in all groups at 2 and 4 hours ($P < 0.05$ versus ST each). Notably, LVSWI was higher in the V₂R-antagonist group than in the AVP

group at 8 and 10 hours (Table 1). Left ventricular contractility, expressed as a Starling-based relationship between LVSWI and preload (pulmonary artery occlusion pressure), was higher in animals treated with the V₂R-antagonist than with placebo (Figure 1b). Cardiac index increased after ST. Heart rate was lower following AVP infusion than in both other groups (8 hours: $P = 0.027$ versus V₂R-antagonist; $P = 0.031$ versus placebo; Table 1).

Central venous and pulmonary artery occlusion pressures, as surrogate variables of cardiac filling pressures, increased in all groups as compared with ST but were higher in animals treated with the V₂R-antagonist as compared with both other groups (Figure 2a,b). Independently

Table 1 Cardiopulmonary variables and mesenteric blood flow

Variable	Group	Baseline	Shock time	4 hours	8 hours	10 hours
HR, beats per min	Placebo	96 ± 2	103 ± 4	123 ± 7 ^a	115 ± 7	102 ± 5
	AVP	93 ± 2	101 ± 5	112 ± 6	99 ± 5 ^b	100 ± 2
	V ₂ antagonist	95 ± 4	102 ± 3	112 ± 6 ^a	115 ± 3 ^{a,c}	101 ± 2
CI, L/min per m ²	Placebo	5.5 ± 0.3	5.8 ± 0.5	8.6 ± 0.8 ^a	7.9 ± 0.5 ^a	5.8 ± 0.6
	AVP	5.2 ± 0.3	6.5 ± 0.4	8.5 ± 0.9	6.4 ± 0.8	5.4 ± 0.8
	V ₂ antagonist	5.3 ± 0.2	5.9 ± 0.3	9.7 ± 0.5 ^a	8.2 ± 0.5 ^a	7.1 ± 0.4
SVRI, dyne-s/cm ⁵ per m ²	Placebo	1,285 ± 109	758 ± 52 ^d	636 ± 60	463 ± 38 ^a	457 ± 107 ^a
	AVP	1,427 ± 101	664 ± 47 ^d	596 ± 109	498 ± 84	479 ± 97 ^a
	V ₂ antagonist	1,406 ± 25	714 ± 46 ^d	509 ± 76 ^a	388 ± 54 ^a	464 ± 75 ^a
MAP, mm Hg	Placebo	91 ± 4	58 ± 4 ^d	66 ± 3 ^a	55 ± 3	44 ± 3 ^a
	AVP	93 ± 2	57 ± 1 ^d	68 ± 2 ^a	56 ± 4	43 ± 1 ^a
	V ₂ antagonist	96 ± 2	58 ± 1 ^d	68 ± 3 ^a	54 ± 2	51 ± 3 ^a
SVI, mL/m ²	Placebo	59 ± 4	58 ± 7	78 ± 3 ^a	68 ± 3	55 ± 7
	AVP	56 ± 3	64 ± 2	80 ± 7	63 ± 7	55 ± 9
	V ₂ antagonist	53 ± 2	57 ± 3	78 ± 7 ^a	71 ± 5	70 ± 3
LVSWI, g/m per m ²	Placebo	67 ± 3	41 ± 4 ^d	64 ± 6 ^a	43 ± 4	22 ± 4 ^a
	AVP	67 ± 3	42 ± 2 ^d	60 ± 3 ^a	26 ± 2 ^b	21 ± 4 ^a
	V ₂ antagonist	65 ± 3	37 ± 2 ^d	65 ± 5 ^a	36 ± 1 ^c	29 ± 2 ^c
MPAP, mm Hg	Placebo	14 ± 1	20 ± 1 ^d	22 ± 1	24 ± 2 ^a	26 ± 2 ^a
	AVP	15 ± 0	18 ± 1 ^d	22 ± 1 ^a	25 ± 1 ^a	27 ± 2 ^a
	V ₂ antagonist	15 ± 1	21 ± 1 ^d	25 ± 2 ^a	27 ± 1 ^a	29 ± 1 ^a
PVRI, dyne-s/cm ⁵ per m ²	Placebo	106 ± 8	139 ± 22	119 ± 15	119 ± 12	144 ± 30
	AVP	124 ± 9	143 ± 8	90 ± 13 ^a	81 ± 16 ^a	150 ± 29
	V ₂ antagonist	129 ± 9	150 ± 9	121 ± 26	103 ± 8 ^a	123 ± 10
Qma, % of baseline	Placebo	100 ± 0	109 ± 17	135 ± 27	94 ± 17	60 ± 10 ^a
	AVP	100 ± 0	95 ± 7	118 ± 21	86 ± 16	41 ± 8 ^a
	V ₂ antagonist	100 ± 0	95 ± 11	115 ± 11	75 ± 6	43 ± 8 ^a

^a*P* < 0.05 versus shock time; ^b*P* < 0.05 versus placebo; ^c*P* < 0.05 versus arginine vasopressin (AVP); ^d*P* < 0.05 versus baseline; each group *n* = 7. CI, cardiac index; HR, heart rate; LVSWI, left ventricular stroke work index; MAP, mean arterial pressure; MPAP, mean pulmonary arterial pressure; PVRI, pulmonary vascular resistance index; Qma, mesenteric arterial blood flow; SVI, stroke volume index; SVRI, systemic.

from the individual treatment regimen, mean pulmonary artery pressure increased during the study period (8 and 10 hours: *P* < 0.05 versus ST each; Table 1).

Mesenteric blood flow

Mesenteric blood flow decreased in all groups (10 hours: *P* < 0.05 versus ST each; Table 1) without any statistically significant differences among groups.

Pulmonary gas exchange and global oxygen transport

Besides a lower PaO₂/FiO₂ (arterial partial pressure of oxygen/fraction of inspired oxygen) ratio in the V₂R-antagonist group compared with the placebo group at 4 hours (*P* = 0.039, Table 2), there were no statistically significant differences between study groups in variables of pulmonary gas exchange and global oxygen transport (Table 2).

Capillary leakage

In all study groups, septic shock was characterized by a marked decrease in plasma protein concentrations

(ST: *P* < 0.001 versus BL each) that progressed over the study period (8 hours: *P* < 0.001 versus ST each; Table 3). At the same time, there were no statistical differences in hematocrit within or among groups (Table 2), suggesting adequate fluid resuscitation. Cumulative positive net fluid balance was similar with all three treatment regimes (V₂R-antagonist: 19 ± 1 mL/kg per hour; AVP: 17 ± 1 mL/kg per hour; placebo: 18 ± 2 mL/kg per hour).

Metabolic changes and electrolytes

Septic shock was associated with decreases in arterial pH and base excess (*P* < 0.05 versus BL each and *P* < 0.001 versus BL each, respectively) and increases in arterial lactate concentrations (*P* < 0.05 versus BL each) in all groups (Figure 3a,b and Table 2). These metabolic changes progressed during the observation period (8 hours: *P* < 0.001 versus ST each). However, the increase in arterial lactate concentration was attenuated (8 and 10 hours: *P* < 0.01 each), arterial base excess was less negative, and pH values were higher in the selective V₂R-antagonist group as compared with the AVP and placebo groups after 8 hours

Table 2 Hematocrit, electrolytes, acid-base balance, and global oxygen transport

Variable	Group	Baseline	Shock time	4 hours	8 hours	10 hours
Hct, %	Placebo	30 ± 2	28 ± 2	30 ± 2	30 ± 2	27 ± 2
	AVP	27 ± 2	26 ± 2	28 ± 2	27 ± 1	28 ± 2
	V ₂ antagonist	26 ± 1	25 ± 2	27 ± 2	26 ± 2	27 ± 1
Na ⁺ , mmol/L	Placebo	141 ± 1	140 ± 1	140 ± 1	140 ± 1	140 ± 1
	AVP	140 ± 1	139 ± 1	139 ± 1	139 ± 1	138 ± 1
	V ₂ antagonist	140 ± 0	139 ± 1	140 ± 1	140 ± 1	140 ± 1
K ⁺ , mmol/L	Placebo	4.1 ± 0.1	4.3 ± 0.2	4.4 ± 0.3	5.5 ± 0.3 ^a	6.1 ± 0.3 ^a
	AVP	3.8 ± 0.2	4.0 ± 0.2	4.1 ± 0.1	5.2 ± 0.3 ^a	5.6 ± 0.4 ^a
	V ₂ antagonist	3.9 ± 0.3	4.2 ± 0.3	4.3 ± 0.2	5.1 ± 0.3	5.5 ± 0.4 ^a
Cl ⁻ , mmol/L	Placebo	108 ± 1	117 ± 2 ^b	120 ± 1	124 ± 1 ^a	125 ± 1 ^a
	AVP	105 ± 1	113 ± 1 ^b	118 ± 1	121 ± 1 ^a	123 ± 1 ^a
	V ₂ antagonist	108 ± 1	115 ± 2 ^b	118 ± 2	121 ± 2	122 ± 2 ^a
pH _a , -log ₁₀ [H ⁺]	Placebo	7.39 ± 0.01	7.30 ± 0.02 ^b	7.20 ± 0.02	7.09 ± 0.04 ^a	7.01 ± 0.06 ^a
	AVP	7.42 ± 0.01	7.31 ± 0.02 ^b	7.22 ± 0.02	7.05 ± 0.05 ^a	7.04 ± 0.06 ^a
	V ₂ antagonist	7.42 ± 0.02	7.33 ± 0.02 ^b	7.28 ± 0.01	7.22 ± 0.04 ^{c,d}	7.11 ± 0.05 ^a
PaO ₂ /FiO ₂ , mm Hg	Placebo	516 ± 23	458 ± 26	435 ± 43	217 ± 41 ^a	149 ± 32 ^a
	AVP	488 ± 23	492 ± 55	383 ± 27 ^a	141 ± 25 ^a	160 ± 19 ^a
	V ₂ antagonist	465 ± 27	412 ± 26	313 ± 20 ^{a,c}	153 ± 30 ^a	140 ± 26 ^a
SvO ₂ , %	Placebo	78 ± 3	74 ± 4	80 ± 3	74 ± 1	60 ± 4 ^a
	AVP	78 ± 1	76 ± 2	83 ± 4	70 ± 5	72 ± 4
	V ₂ antagonist	79 ± 2	78 ± 2	85 ± 2	78 ± 3	68 ± 4
DO ₂ l, mL/min per m ²	Placebo	731 ± 63	719 ± 83	1,105 ± 115 ^a	918 ± 39	575 ± 92
	AVP	641 ± 58	739 ± 65	955 ± 128	749 ± 99	620 ± 85
	V ₂ antagonist	598 ± 36	664 ± 50	1,132 ± 139 ^a	936 ± 50	707 ± 64
VO ₂ l, mL/min per m ²	Placebo	160 ± 12	179 ± 14	181 ± 19	172 ± 22	155 ± 21
	AVP	163 ± 13	167 ± 8	175 ± 8	144 ± 25	123 ± 18 ^a
	V ₂ antagonist	128 ± 17	153 ± 10	163 ± 17	142 ± 13	132 ± 17
O ₂ -ER, %	Placebo	23 ± 3	26 ± 3	18 ± 3	18 ± 2 ^a	26 ± 2
	AVP	24 ± 2	25 ± 1	21 ± 6	21 ± 4	22 ± 4
	V ₂ antagonist	20 ± 1	23 ± 1	13 ± 1 ^a	16 ± 2	20 ± 4

^a*P* < 0.05 versus shock time; ^b*P* < 0.05 versus baseline; ^c*P* < 0.05 versus placebo; ^d*P* < 0.05 versus arginine vasopressin; each group *n* = 7. AVP, arginine vasopressin; DO₂l, oxygen delivery index; Hct, hematocrit; O₂-ER, oxygen extraction rate; PaO₂/FiO₂, ratio of arterial partial pressure of oxygen and inspiratory oxygen fraction; pH_a, arterial potentia hydrogenii; SvO₂, mixed venous oxygen saturation, VO₂l, oxygen consumption index.

(*P* < 0.05 each). Plasma concentrations of potassium and chloride increased in all groups during the study period (*P* < 0.05 versus ST each) without significant differences among groups.

Laboratory variables of organ function and arginine vasopressin plasma levels

Alanine aminotransferase and aspartate aminotransferase activity as well as plasma concentrations of bilirubin were reduced by the selective V₂R-antagonist as compared with placebo animals (8 hours: *P* < 0.05 each; Table 3). Renal dysfunction was evidenced by a progressive increase in blood urea nitrogen and plasma creatinine concentrations as well as a decrease in urine output and creatinine clearance in placebo animals (Figure 4 and Table 3). Infusion of the selective V₂R-antagonist was associated with an increased creatinine clearance (4 hours: *P* < 0.001), a higher urine output (2 to 4 hours: *P* < 0.001 each), and lower

blood urea nitrogen levels (4 to 8 hours: *P* = 0.031 and *P* = 0.023, respectively) as compared with the placebo group. There were no statistical differences in renal and liver function between the V₂R-antagonist and the AVP group.

The onset of septic shock was associated with an increase in AVP plasma levels as compared with BL in all groups (*P* < 0.05 versus BL each; Figure 5). Whereas AVP plasma levels remained constant in the placebo group, infusion of AVP increased AVP plasma levels up to 149 ± 21 pg/mL. Treatment with the selective V₂R-antagonist led to a significant decrease of AVP plasma levels as compared with ST (*P* < 0.001) and with both other groups (4 to 8 hours: *P* < 0.05 versus placebo; *P* < 0.001 versus AVP).

Immunohistochemical analyses

Immunohistochemical analyses of lung tissue revealed an increase in hemeoxygenase-1 concentration in the selective V₂R-antagonist group as compared with

Table 3 Surrogate parameters of organ (dys)function

Variable	Group	Baseline	Shock time	4 hours	8 hours
AST, U/L	Placebo	71 ± 7	76 ± 6	81 ± 14	112 ± 18 ^a
	AVP	71 ± 7	78 ± 7	80 ± 12	77 ± 8
	V ₂ antagonist	72 ± 8	74 ± 8	58 ± 9	63 ± 10 ^b
ALT, U/L	Placebo	7 ± 2	9 ± 1	9 ± 2	13 ± 3
	AVP	8 ± 3	11 ± 1	8 ± 2	11 ± 2
	V ₂ antagonist	8 ± 2	10 ± 3	5 ± 1	6 ± 1 ^b
Bilirubin, mg/dL	Placebo	0.24 ± 0.02	0.24 ± 0.02	0.26 ± 0.04	0.25 ± 0.02
	AVP	0.25 ± 0.02	0.23 ± 0.02	0.23 ± 0.02	0.18 ± 0.04
	V ₂ antagonist	0.24 ± 0.02	0.23 ± 0.02	0.23 ± 0.03	0.16 ± 0.03 ^b
Plasma protein, mg/dL	Placebo	4.3 ± 0.2	1.9 ± 0.2 ^c	1.2 ± 0.1 ^a	0.7 ± 0.0 ^a
	AVP	4.4 ± 0.2	2.1 ± 0.1 ^c	1.2 ± 0.1 ^a	0.9 ± 0.2 ^a
	V ₂ antagonist	4.2 ± 0.3	1.9 ± 0.2 ^c	1.2 ± 0.1	0.9 ± 0.2 ^a
Creatinine, mg/dL	Placebo	0.8 ± 0.1	0.7 ± 0.1	1.1 ± 0.1	1.5 ± 0.1 ^a
	AVP	0.7 ± 0.1	0.7 ± 0.1	0.7 ± 0.1	1.3 ± 0.2 ^a
	V ₂ antagonist	0.8 ± 0.1	0.8 ± 0.1	0.8 ± 0.1	1.1 ± 0.2
Creatinine clearance, mL/min	Placebo	270 ± 82	228 ± 36	37 ± 10 ^a	16 ± 3 ^a
	AVP	254 ± 29	197 ± 42	214 ± 59 ^b	24 ± 2 ^a
	V ₂ antagonist	235 ± 43	198 ± 20	346 ± 52 ^b	48 ± 15 ^a

^a*P* < 0.05 versus shock time; ^b*P* < 0.05 versus placebo; ^c*P* < 0.05 versus baseline; *n* = 7 each. ALT, alanine aminotransferase; AST, aspartate aminotransferase; AVP, arginine vasopressin.

placebo animals (*P* = 0.047; Figure 6a). In addition, pulmonary 3-nitrotyrosine concentrations were lower in animals treated with the selective V₂R-antagonist as compared with AVP (*P* = 0.017; *P* = 0.056 versus placebo; Figure 6b).

Survival time

All animals died within 17 hours after the onset of septic shock (Figure 7). Sheep treated with the selective V₂R-antagonist had a longer survival time (14 ± 1 hours) than animals that received AVP (11 ± 1 hours; *P* = 0.007) or

placebo (11 ± 1 hours; *P* = 0.025). There were no significant differences in survival time between the AVP and sole norepinephrine groups (*P* = 0.727).

Discussion

The major findings of the present study are that first-line therapy with the selective V₂R-antagonist (a) stabilized cardiopulmonary hemodynamics as effectively, (b) increased cardiac filling pressures, (c) attenuated metabolic acidosis, (d) limited myocardial and renal dysfunction, (e) reduced AVP plasma levels, (f) attenuated tissue injury

secondary to nitrosative stress, and (g) slightly prolonged survival in early volume-resuscitated, hyperdynamic ovine septic shock when compared with placebo and AVP infusion.

The relative vasopressin deficiency [18] represents the rationale for the use of AVP in the treatment of septic shock. However, only one third of septic shock patients suffer from low AVP plasma levels [19]. Typically, endogenous AVP secretion increases in the early phase of septic shock and decreases thereafter. Since V₂Rs are involved in several characteristic pathways of septic shock [4-11,20], selective V₂R-antagonism rather than V₂R-stimulation (for example, via AVP infusion) may be advantageous under these circumstances.

In the present study, AVP plasma levels increased with the onset of septic shock in all groups and remained at

this level in the placebo group during the whole study period. The absence of a 'relative vasopressin deficiency' may be one reason for the ineffectiveness of AVP in reducing norepinephrine requirements as compared with standard treatment with norepinephrine in the placebo group. Another potential explanation is that the AVP dose of 0.05 µg/kg per hour (equivalent to 0.5 mU/kg per minute or 0.035 U/minute in a 70-kg patient) might have been insufficient for the fulminant injury in our model (100% mortality within 17 hours). The latter assumption is in harmony with the observation made in VASST that AVP reduced mortality in less severe septic shock but not in the more severe septic shock population [2]. In this context, Torgersen and colleagues [21] recently reported that, in patients with sepsis-induced vasodilatory shock, a supplementary infusion of 0.067 U/minute AVP was more effective in restoring MAP and reducing norepinephrine requirements than the recommended low dose of 0.033 U/minute.

Interestingly, infusion of the selective V₂R-antagonist reduced AVP plasma levels as compared with AVP- and placebo-treated animals. This finding appears to be surprising at first glance. In this context, however, it may be of importance that AVP has a positive feedback on its own release via V₂R [22]. Therefore, it is most likely that inhibition of this mechanism has accounted for the low AVP plasma levels noticed in the V₂R-antagonist group.

Another interesting result of the present study is that the selective V₂R-antagonist was as effective as AVP in stabilizing cardiopulmonary hemodynamics without increasing volume and norepinephrine requirements. The reduction in metabolic acidosis by the V₂R-antagonist - as suggested by higher pH values, less negative base excess, and lower lactate levels as compared with both other

Figure 6 Pulmonary hemoxygenase-1 (a) and 3-nitrotyrosine (b) concentrations. [†]*P* < 0.05 versus placebo; [§]*P* < 0.05 versus arginine vasopressin (AVP); *n* = 7 each. 3-NT, 3-nitrotyrosine; HO-1, hemoxygenase-1.

groups - probably reduced systemic vasodilation [23] and contributed to an improved efficacy of norepinephrine by increasing the adrenergic receptor sensitivity [24,25].

In this context, it may also be important that extrarenal V₂R mediates vasorelaxant effects [4], thereby decreasing MAP and vascular resistance not only in the experimental setting [26] but also in humans [6,27].

In addition, the increased cardiac filling pressures in animals treated with the V₂R-antagonist may have improved systemic hemodynamics. This assumption is supported by the Starling-based relationship between LVSWI and preload (Figure 1b). Since hematocrit remained stable in all groups, the increased preload in

the V₂R-antagonist group has most likely been caused by a mobilization of fluid from venous capacity vessels.

Whereas both the V₂R-antagonist and AVP increased urine output and creatinine clearance as compared with placebo animals, the V₂R-antagonist additionally reduced blood urea nitrogen versus placebo. A protective effect of V₂R-antagonism on renal function is supported by Rondaij and colleagues [28], who reported that V₂R agonism caused histological renal lesions in rats and that these lesions were prevented by V₂R-antagonism.

In addition, the reduction of oxidative stress, as suggested by immunohistochemical analyses of lung tissues, probably contributed to the attenuated organ dysfunction in the V₂R-antagonist group as compared with placebo and AVP. Whereas 3-nitrotyrosine represents a stable *in vivo* biomarker of the highly cytotoxic compound peroxynitrite [29], hemoxygenase-1 has been reported to provide cytoprotective effects [30].

Attenuation of cardiovascular, metabolic, and renal function as well as nitrosative stress in response to first-line V₂R-antagonist infusion led to a slight prolongation in survival time as compared with AVP and placebo treatment. Such effects on survival time were not observed with AVP, suggesting that its V₂R agonism might potentially be disadvantageous.

This study has some limitations that we want to acknowledge. In the absence of source control and antibiotic therapy, the present model was associated with a high mortality (all animals died within the observation period). As a consequence, effects of the investigated therapeutic approaches could be analyzed only during the acute phase of the injury. In addition, the present study was not designed primarily for detecting differences in mortality. For these reasons, data on survival times in

Figure 7 Kaplan-Meier survival curve. [†]*P* < 0.05 versus placebo; [§]*P* < 0.05 versus arginine vasopressin (AVP); *n* = 7 each. ST, shock time.

the current study should not be overestimated. In addition, conclusions on the clinical relevance of the present findings are limited by the experimental design and the use of previously healthy animals, whereas the majority of patients typically suffer from comorbidities. Finally, the risk of false-positive results in a study with numerous outcome variables and time points has to be taken into consideration.

Conclusions

To our knowledge, this is the first study providing evidence that, under conditions with high endogenous AVP plasma levels, first-line treatment with the selective V₂R-antagonist supplemented with open-label norepinephrine improves cardiovascular, metabolic, liver, and renal function and slightly prolongs survival when compared with first-line therapy with AVP or placebo in ovine septic shock. On the basis of the present findings, the use of selective V₂R-antagonists potentially represents a new therapeutic approach in the early stage of septic shock.

Key messages

- V₂-receptor stimulation aggravates sepsis-induced vasodilation, fluid accumulation, and microvascular thrombosis.
- Arginine vasopressin (AVP) infusion in septic shock may be less effective when endogenous AVP plasma levels are high.
- In ovine septic shock, selective V₂-receptor-antagonism supplemented with open-label norepinephrine stabilized cardiovascular hemodynamics as effectively as combined AVP and open-label norepinephrine.
- Selective V₂-receptor-antagonism attenuated metabolic, liver, and renal dysfunction as compared with AVP and placebo therapy in ovine septic shock.
- Selective V₂-receptor-antagonism might represent a useful therapeutic option in septic shock under conditions with high endogenous AVP plasma levels.

Additional material

Additional file 1: Supplemental Digital Content. Additional information on the methods and procedures applied in the present study [31-33].

Abbreviations

AVP: arginine vasopressin; BL: baseline; ELISA: enzyme-linked immunosorbent assay; LVSWI: left ventricular stroke work index; MAP: mean arterial pressure; ST: shock time; V_{1a}R/V₂R: V_{1a}/V₂ receptor; VASST: Vasopressin and Septic Shock Trial.

Acknowledgements

The authors thank Mareike Schneider, a medical student from the Department of Anesthesiology and Intensive Care at the University of Muenster (Muenster, Germany), for expert technical assistance during the

study. This work was supported only by intramural funding of the University of Muenster.

Author details

¹Department of Anesthesiology and Intensive Care, University of Muenster, Albert-Schweitzer-Str. 33, Muenster 48149, Germany. ²Department of Anesthesiology and Intensive Care, University of Rome, 'La Sapienza', Viale del Policlinico 155, 00161 Rome, Italy. ³Department of Biostatistics and Epidemiology, University of Texas Medical Branch, 301 University Boulevard, Galveston, TX 77550, USA. ⁴Department of Intensive Care Medicine, Inselspital, Medical University of Bern, CH-3010 Bern, Switzerland. ⁵Investigational Intensive Care Unit, Department of Anesthesiology, University of Texas Medical Branch, 301 University Boulevard, Galveston, TX 77550, USA.

Authors' contributions

SR designed and performed the experiment, summarized and analyzed the data, and wrote the manuscript. CE designed and performed the experiment, summarized and analyzed the data, and edited the manuscript. MW and AM designed the experiment, analyzed the data, and edited the manuscript. ML, EW, MD, HVA, and DLT analyzed the data and edited the manuscript. A-KS, EL, and TKG performed the experiment and summarized the data. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 30 April 2010 Revised: 14 June 2010

Accepted: 5 November 2010 Published: 5 November 2010

References

1. Dellinger RP, Levy MM, Carlet JM, Bion J, Parker MM, Jaeschke R, Reinhart K, Angus DC, Brun-Buisson C, Beale R, Calandra T, Dhainaut JF, Gerlach H, Harvey M, Marini JJ, Marshall J, Ranieri M, Ramsay G, Sevransky J, Thompson BT, Townsend S, Vender JS, Zimmerman JL, Vincent JL: **Surviving Sepsis Campaign: international guidelines for management of severe sepsis and septic shock: 2008.** *Crit Care Med* 2008, **36**:296-327.
2. Russell J, Walley K, Singer J, Gordon A, Hébert P, Cooper D, Holmes C, Mehta S, Granton J, Storms M, Cook D, Presneill J, Ayers D: **Vasopressin versus Norepinephrine Infusion in Patients with Septic Shock.** *N Engl J Med* 2008, **358**:877-887.
3. Petersen MB: **The effect of vasopressin and related compounds at V1a and V2 receptors in animal models relevant to human disease.** *Basic Clin Pharmacol Toxicol* 2006, **99**:96-103.
4. Barrett LK, Singer M, Clapp LH: **Vasopressin: mechanisms of action on the vasculature in health and in septic shock.** *Crit Care Med* 2007, **35**:33-40.
5. Kaufmann JE, Vischer UM: **Cellular mechanisms of the hemostatic effects of desmopressin (DDAVP).** *J Thromb Haemost* 2003, **1**:682-689.
6. Bichet DG, Razi M, Loneragan M, Arthus MF, Papukna V, Kortas C, Barjon JN: **Hemodynamic and coagulation responses to 1-desamino[8-D-arginine] vasopressin in patients with congenital nephrogenic diabetes insipidus.** *N Engl J Med* 1988, **318**:881-887.
7. Liard JF: **cAMP and extrarenal vasopressin V2 receptors in dogs.** *Am J Physiol* 1992, **263**:H1888-1891.
8. Kaufmann JE, Lezzi M, Vischer UM: **Desmopressin (DDAVP) induces NO production in human endothelial cells via V2 receptor- and cAMP-mediated signaling.** *J Thromb Haemost* 2003, **1**:821-828.
9. Traber DL: **Selective V1a receptor agonists in experimental septic shock [Abstract].** *Crit Care* 2007, **11**:P51.
10. Kanwar S, Woodman RC, Poon MC, Murohara T, Lefer AM, Davenpeck KL, Kubes P: **Desmopressin induces endothelial P-selectin expression and leukocyte rolling in postcapillary venules.** *Blood* 1995, **86**:2760-2766.
11. Rehberg S, Laporte R, Enkhbaatar P, La E, Wisniewski K, Traber LD, Rivière P, Traber DL: **Arginine vasopressin increases plasma levels of von Willebrand factor in sheep.** *Crit Care* 2009, **13**:P182.
12. Manning M, Klis WA, Kruszynski M, Przybylski JP, Olma A, Wo NC, Pelton GH, Sawyer WH: **Novel linear antagonists of the antidiuretic (V2) and vasopressor (V1) responses to vasopressin.** *Int J Pept Protein Res* 1988, **32**:455-467.
13. Manning M, Przybylski J, Grzonka Z, Nawrocka E, Lammek B, Misicka A, Cheng LL, Chan WY, Wo NC, Sawyer WH: **Potent V2/V1a vasopressin**

- antagonists with C-terminal ethylenediamine-linked retro-amino acids. *J Med Chem* 1992, **35**:3895-3904.
14. Rehberg S, Ertmer C, Kohler G, Spiegel HU, Morelli A, Lange M, Moll K, Schlack K, Van Aken H, Su F, Vincent JL, Westphal M: **Role of arginine vasopressin and terlipressin as first-line vasopressor agents in fulminant ovine septic shock.** *Intensive Care Med* 2009, **35**:1286-1296.
 15. Wang Z, Su F, Rogiers P, Vincent JL: **Beneficial effects of recombinant human activated protein C in a ewe model of septic shock.** *Crit Care Med* 2007, **35**:2594-2600.
 16. Westphal M, Enkhbaatar P, Schmalstieg FC, Kulp GA, Traber LD, Morita N, Cox RA, Hawkins HK, Westphal-Varghese BB, Rudloff HE, Maybauer DM, Maybauer MO, Burke AS, Murakami K, Saunders F, Horvath EM, Szabo C, Traber DL: **Neuronal nitric oxide synthase inhibition attenuates cardiopulmonary dysfunctions after combined burn and smoke inhalation injury in sheep.** *Crit Care Med* 2008, **36**:1196-1204.
 17. Westphal M, Cox RA, Traber LD, Morita N, Enkhbaatar P, Schmalstieg FC, Hawkins HK, Maybauer DM, Maybauer MO, Murakami K, Burke AS, Westphal-Varghese BB, Rudloff HE, Salisbury JR, Jodoin JM, Lee S, Traber DL: **Combined burn and smoke inhalation injury impairs ovine hypoxic pulmonary vasoconstriction.** *Crit Care Med* 2006, **34**:1428-1436.
 18. Landry DW, Levin HR, Gallant EM, Ashton RC Jr, Seo S, D'Alessandro D, Oz MC, Oliver JA: **Vasopressin deficiency contributes to the vasodilation of septic shock.** *Circulation* 1997, **95**:1122-1125.
 19. Sharshar T, Blanchard A, Paillard M, Raphael JC, Gajdos P, Annane D: **Circulating vasopressin levels in septic shock.** *Crit Care Med* 2003, **31**:1752-1758.
 20. Rehberg S, Ertmer C, Traber DL, Van Aken H, Westphal M: **Selective V2-receptor-antagonism as a new therapeutic approach in ovine septic shock.** *Intensive Care Med* 2009, **35**:111, #422.
 21. Torgersen C, Dünser MW, Wenzel V, Jochberger S, Mayr V, Schmittinger CA, Lorenz I, Schmid S, Westphal M, Grander W, Luckner G: **Comparing two different arginine vasopressin doses in advanced vasodilatory shock: a randomized, controlled, open-label trial.** *Intensive Care Med* 2010, **36**:57-65.
 22. Landgraf R, Ramirez AD, Ramirez VD: **The positive feedback action of vasopressin on its own release from rat septal tissue in vitro is receptor-mediated.** *Brain Res* 1991, **545**:137-141.
 23. Pedoto A, Caruso JE, Nandi J, Oler A, Hoffmann SP, Tassiopoulos AK, McGraw DJ, Camporesi EM, Hakim TS: **Acidosis stimulates nitric oxide production and lung damage in rats.** *Am J Respir Crit Care Med* 1999, **159**:397-402.
 24. Simonis G, Marquetant R, Rothele J, Strasser RH: **The cardiac adrenergic system in ischaemia: differential role of acidosis and energy depletion.** *Cardiovasc Res* 1998, **38**:646-654.
 25. Ryan AJ, Gisolfi CV: **Responses of rat mesenteric arteries to norepinephrine during exposure to heat stress and acidosis.** *J Appl Physiol* 1995, **78**:38-45.
 26. Liard JF: **Interaction between V1 and V2 effects in hemodynamic response to vasopressin in dogs.** *Am J Physiol* 1990, **258**:H482-489.
 27. Tagawa T, Imaizumi T, Shiramoto M, Endo T, Hironaga K, Takeshita A: **V2 receptor-mediated vasodilation in healthy humans.** *J Cardiovasc Pharmacol* 1995, **25**:387-392.
 28. Rondajj MG, Bierings R, Kragt A, Gijzen KA, Sellink E, van Mourik JA, Fernandez-Borja M, Voorberg J: **Dynein-dynactin complex mediates protein kinase A-dependent clustering of Weibel-Palade bodies in endothelial cells.** *Arterioscler Thromb Vasc Biol* 2006, **26**:49-55.
 29. Radi R: **Nitric oxide, oxidants, and protein tyrosine nitration.** *Proc Natl Acad Sci USA* 2004, **101**:4003-4008.
 30. Chung SW, Liu X, Macias AA, Baron RM, Perrella MA: **Heme oxygenase-1-derived carbon monoxide enhances the host defense response to microbial sepsis in mice.** *J Clin Invest* 2008, **118**:239-247.
 31. Su F, Wang Z, Cai Y, Rogiers P, Vincent JL: **Fluid resuscitation in severe sepsis and septic shock: albumin, hydroxyethyl starch, gelatin or ringer's lactate-does it really make a difference?** *Shock* 2007, **27**:520-526.
 32. **Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. The Acute Respiratory Distress Syndrome Network.** *N Engl J Med* 2000, **342**:1301-1308.
 33. Westphal M, Stubbe H, Sielenkamper AW, Ball C, Van Aken H, Borgulya R, Bone HG: **Effects of titrated arginine vasopressin on hemodynamic variables and oxygen transport in healthy and endotoxemic sheep.** *Crit Care Med* 2003, **31**:1502-1508.

doi:10.1186/cc9320

Cite this article as: Rehberg et al.: Role of selective V₂-receptor-antagonism in septic shock: a randomized, controlled, experimental study. *Critical Care* 2010 **14**:R200.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit