

LETTER

Search for biomarkers in critically ill patients: a new approach based on nuclear magnetic resonance spectroscopy of mini-bronchoalveolar lavage fluid

Chandan Singh^{1,2}, Ratan Kumar Rai¹, Afzal Azim^{3*}, Neeraj Sinha^{1*} and Arvind Kumar Baronia³

Human lungs have the function of gas exchange in the body, performed efficiently by the unique anatomy of the alveoli. The alveolar epithelial lining fluid, reflecting a snapshot of molecular events happening there, can be extracted by bronchoscopic/nonbronchoscopic methods and used to study critical diseases. Although there are methods to study the pathophysiological conditions, there is still a need for newer and faster methods that can provide metabolic information about disease diagnosis, severity and progression.

In this letter we present nuclear magnetic resonance-based metabolomics (a key component of system biology), which has potential for disease diagnosis and treatment monitoring [1].

Acute respiratory distress syndrome (ARDS) is a disease with a high rate of mortality and morbidity worldwide, survival being only up to 40%. There is immense need for biomarkers associated with ARDS, and scientists have been working hard for the last four decades to discover these, without the anticipated success. Bronchoalveolar lavage fluid, mini-bronchoalveolar lavage (mBAL) fluid and serum have been the primary body fluids studied for this purpose. Recently, our group explored small molecular weight metabolites responsible for severity of ARDS, employing metabolomics in mBAL fluid and serum [2,3]. Both bronchoalveolar lavage fluid and mBAL fluid can also be used for nuclear magnetic resonance-based metabolomics [4].

We used a nonbronchoscopic, catheter inside catheter technique to extract mBAL fluid [5]. Most of the metabolites were characterized and identified (Figure 1). The branch-chain amino acid, lactate, alanine, lysine, arginine, acetate, succinate, taurine, phenylalanine, betaine and aspartate levels were elevated in the mBAL fluid collected from a diseased patient compared with that from a healthy control (Figure 1). The proline level was found to decrease in the case of ARDS. The roles of the above-mentioned small molecular weight metabolites have been discussed previously [2]. One-dimensional nuclear magnetic resonance spectra can be preprocessed and utilized for unsupervised and supervised chemometric analysis, highlighting the role of key metabolites. Jelly should be avoided during extraction of mBAL fluid because resonance from it dominates the spectrum and masks resonance from small molecular weight metabolites, as shown in Figure 1. We have summarized the complete procedure in Figure 2 and in detail in one of our earlier studies [2].

With precautions, researchers will find application for this procedure in the study of diseases such as respiratory failure, interstitial lung disease, sarcoidosis and so forth. Besides the above-mentioned studies, this will be of extreme importance for clinicians as well as basic scientists trying to obtain more information about diseases where the balance of bronchoalveolar lavage fluid is affected.

* Correspondence: draazim2002@gmail.com; neeraj.sinha@cbmr.res.in

³Department of Critical Care Medicine, Sanjay Gandhi Postgraduate Institute of Medical Sciences, Lucknow 226014, India

¹Centre of Biomedical Research, Sanjay Gandhi Postgraduate Institute of Medical Sciences Campus, Raibareilly Road, Lucknow 226014, India

Full list of author information is available at the end of the article




Figure 1 Interference due to jelly. (a) Spectrum showing resonance from jelly. (b) Spectrum of mini-bronchoalveolar lavage (mBAL) fluid dominated by resonance from jelly. (c) Spectrum showing resonance originating from small metabolites present inside mBAL fluid (control). (d) Various small molecular weight metabolites present inside mBAL fluid of an acute respiratory distress syndrome patient. 1, isoleucine, valine and leucine; 2, ethanol; 3, lactate/threonine; 4, alanine; 5, arginine and lysine; 6, acetate; 7, glutamate; 8, succinate; 9, pyruate; 10, glutamine; 11, aspartate; 12, asparagine; 13, creatine and lysine; 14, histidine; 15, betaine; 16, taurine; 17, choline; 18, glycine; 19, lactate; 20, threonine; 21, β -glucose; 22, uracil/urea; 23, fumaric acid; 24 and 26, tyrosine; 25 and 29, histidine; 27, phenylalanine; 28, uracil; 30, adenine; 31, formate.




Figure 2 Complete process of extracting and processing mini-bronchoalveolar lavage fluid and recording its nuclear magnetic resonance spectrum. 1D, one-dimensional; 2D, two-dimensional; ARDS, acute respiratory distress syndrome; mBAL, mini-bronchoalveolar lavage; NMR, nuclear magnetic resonance; PBS, phosphate-buffered saline.

Authors' contributions

CS and RKR carried out sample collection, performed nuclear magnetic resonance analysis and wrote the manuscript. NS and AA designed the work. AA and AKB performed extraction of mBAL fluid from the patients. All authors read and approved the final version of the manuscript.

Acknowledgements

CS and RKR acknowledge financial support as a fellowship from the Council of Scientific and Industrial Research, India. This work was financially supported by the Department of Biotechnology, India (Grant Number BT/PR12700/BRB/10/719/2009) through which important consumables were purchased. The funding agency had no role in the design of study.

Author details

¹Centre of Biomedical Research, Sanjay Gandhi Postgraduate Institute of Medical Sciences Campus, Raibareilly Road, Lucknow 226014, India. ²School of Biotechnology, Faculty of Science, Banaras Hindu University, Varanasi 221005, India. ³Department of Critical Care Medicine, Sanjay Gandhi Postgraduate Institute of Medical Sciences, Lucknow 226014, India.

Published online: 20 November 2014

References

- Beckonert O, Keun HC, Ebbels TMD, Bundy J, Holmes E, Lindon JC, Nicholson JK: **Metabolic profiling, metabolomic and metabonomic procedures for NMR spectroscopy of urine, plasma, serum and tissue extracts.** *Nat Protoc* 2007, **2**:2692–2703.
- Rai RK, Azim A, Sinha N, Sahoo J, Singh C, Ahmed A, Saigal S, Baronia A, Gupta D, Gurjar M, Poddar B, Singh RK: **Metabolic profiling in human lung injuries by high-resolution nuclear magnetic resonance spectroscopy of bronchoalveolar lavage fluid (BALF).** *Metabolomics* 2013, **9**:667–676.
- Singh C, Rai R, Azim A, Sinha N, Ahmed A, Singh K, Kayastha A, Baronia AK, Gurjar M, Poddar B, Singh R: **Metabolic profiling of human lung injury by ¹H high-resolution nuclear magnetic resonance spectroscopy of blood serum.** *Metabolomics* 2014. [Epub ahead of print].
- Singh C, Rai R, Azim A, Sinha N, Baronia A: **Mini-bronchoalveolar lavage fluid can be used for biomarker identification in patients with lung injury by employing ¹H NMR spectroscopy.** *Crit Care* 2013, **17**:430.
- Khilnani GC, Arafath KT, Hadda V, Kapil A, Sood S, Sharma SK: **Comparison of bronchoscopic and non-bronchoscopic techniques for diagnosis of ventilator associated pneumonia.** *Indian J Crit Care Med* 2011, **15**:16–23.

doi:10.1186/s13054-014-0594-x

Cite this article as: Singh et al.: Search for biomarkers in critically ill patients: a new approach based on nuclear magnetic resonance spectroscopy of mini-bronchoalveolar lavage fluid. *Critical Care* 2014 **18**:594.

Abbreviations

ARDS: acute respiratory distress syndrome; mBAL: mini-bronchoalveolar lavage.

Competing interests

The authors declare that they have no competing interests.