

LETTER

Open Access



Faulty risk-of-bias assessment in a meta-analysis of hydroxyethyl starch for nonseptic ICU patients

Ole Bayer^{1*} and Konrad Reinhart²

See related Research by He et al., <http://www.ccforum.com/content/19/1/92>

Evaluating 6 % hydroxyethyl starch (HES) of different molecular weight and substitution versus other fluids for non-septic patients, He et al. [1] did not confirm the increased mortality, renal replacement therapy (RRT), bleeding and red blood cell transfusions seen in sepsis. A favourable safety profile for 6 % HES solutions was suggested.

He et al. stated that Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines were followed, yet the meta-analysis lacked tests of interaction to compare different HES solutions and pre-specified subgroup analyses. The discussed limitations of the meta-analysis included study quality, heterogeneity, and problematic use [2] of gelatin as a control fluid. The Cochrane Collaboration Risk-of-Bias Tool was cited for guidance in the assessment of study quality, where any use of scales for risk of bias is explicitly

discouraged because summary scores have been shown to be unreliable. He et al. calculated summary scores using a nonvalidated modified Jadad scale [3] where the highest possible score is 8, but the highest possible summary score for He et al. was only 7; reasons for this difference are obscure. Information on how assessment of risk of bias was actually performed is missing. For selective outcome reporting, the study by James et al. was wrongly judged as 'low risk' although 'high risk' had been identified previously [4, 5].

The key message from He et al. that 6 % HES does not seem to increase mortality or RRT incidence in nonseptic ICU patients is speculative because assessment of the study quality is insufficient. HES should be considered unsafe in all ICU patients until proven contrary in high-quality studies.

Authors' response

Bin He, Bo Xu, Xiaoxing Xu, Lixia Li, Rongrong Ren, Zhiyu Chen, Jian Xiao, Yingwei Wang and Bin Xu

We thank Dr Bayer and Dr Reinhart for their comments, which we would like to nuance.

Recently, Myburgh et al. [6] reported that there was no significant difference in 90-day mortality between patients resuscitated with 6 % HES or saline. Another international multicentre large-sample randomised controlled trial (RCT) also reported the use of colloids (including HES, gelatin, dextran and albumin) versus any crystalloids for volume expansion therapy to decrease 90-day mortality [7]. In addition, two meta-analyses also showed that HES did not increase mortality or renal injury in subgroup analyses of nonseptic patients [8, 9]. Our meta-analysis also obtained similar results.

We had pre-specified subgroup analyses by different molecular weight of 6 % HES, risk of bias assessment and different kinds of fluids, all of which were found with no significant difference in mortality. We also made a brief analysis about the limitations of the meta-analysis. The Cochrane Collaboration Risk-of-Bias Tool and the modified Jadad scale were both always used for assessing the risk of bias [3, 8]. We apologise for the clerical error regarding the highest possible score in the modified Jadad scale and the inappropriate judgement for selective outcome reporting of the study by James et al.

Owing to the limitation of the current RCTs, our conclusion did not suggest using HES for these patients, and advised further RCTs of high quality for further study.

* Correspondence: Ole.Bayer@med.uni-jena.de

¹Department of Anaesthesiology and Intensive Care, Jena University Hospital, Erlanger Allee 101, 07747 Jena, Germany

Full list of author information is available at the end of the article

Abbreviations

HES: Hydroxyethyl starch; PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analyses; RCT: Randomised controlled trial; RRT: Renal replacement therapy.

Competing interests

OB received speaker's fees from CSL Behring, Germany. KR received an unrestricted research grant to conduct the VISEP trial and consultancy fees from B Braun Melsungen. The remaining authors declare that they have no competing interests.

Authors' contributions

KR and OB drafted the manuscript. Both authors read and approved the final manuscript.

Author details

¹Department of Anaesthesiology and Intensive Care, Jena University Hospital, Erlanger Allee 101, 07747 Jena, Germany. ²Department of Anaesthesiology and Intensive Care, Center for Sepsis Control and Care, Chairman Global Sepsis Alliance, Jena University Hospital, Erlanger Allee 101, 07747 Jena, Germany.

Published online: 08 October 2015

References

1. He B, Xu B, Xu X, Li L, Ren R, Chen Z, et al. Hydroxyethyl starch versus other fluids for non-septic patients in the intensive care unit: a meta-analysis of randomized controlled trials. *Crit Care*. 2015;19:92.
2. Thomas-Rueddel DO, Vlasakov V, Reinhart K, Jaeschke R, Rueddel H, Hutagalung R, et al. Safety of gelatin for volume resuscitation—a systematic review and meta-analysis. *Intensive Care Med*. 2012;38:1134–42.
3. Bañares R, Albillos A, Rincón D, Alonso S, González M, Ruiz-del-Arbol L, et al. Endoscopic treatment versus endoscopic plus pharmacologic treatment for acute variceal bleeding: a meta-analysis. *Hepatology*. 2002;35:609–15.
4. Finfer S. Hydroxyethyl starch in patients with trauma. *Br J Anaesth*. 2012;108:159–60.
5. Reinhart K, Hartog CS. Hydroxyethyl starch in patients with trauma. *Br J Anaesth*. 2012;108:321–2.
6. Myburgh JA, Finfer S, Bellomo R, Billot L, Cass A, Gattas D, et al. Hydroxyethyl starch or saline for fluid resuscitation in intensive care. *N Engl J Med*. 2012;367:1901–11.
7. Annane D, Siami S, Jaber S, Martin C, Elatrous S, Declère AD, et al. Effects of fluid resuscitation with colloids vs crystalloids on mortality in critically ill patients presenting with hypovolemic shock: the CRISTAL randomized trial. *JAMA*. 2013;310:1809–17.
8. Zarychanski R, Abou-Setta AM, Turgeon AF, Houston BL, McIntyre L, Marshall JC, et al. Association of hydroxyethyl starch administration with mortality and acute kidney injury in critically ill patients requiring volume resuscitation: a systematic review and meta-analysis. *JAMA*. 2013;309:678–88.
9. Muttter TC, Ruth CA, Dart AB. Hydroxyethyl starch (HES) versus other fluid therapies: effects on kidney function. *Cochrane Database Syst Rev*. 2013;7:CD007594.