

ERRATUM

Open Access



Erratum to: Probiotic and synbiotic therapy in critical illness: a systematic review and meta-analysis

William Manzanares¹ and Paul E. Wischmeyer^{2*}

Erratum

Unfortunately, the original version of this article [1] published in *Critical Care* contained an error. We have made an error in the calculation of the pooled risk ratio (RR) and 95% confidence interval (CI) for the effect of probiotics and synbiotics on hospital mortality. Currently, the revised effect on overall mortality is 1.02 (95% CI 0.85,1.22; $P = 0.83$, $I^2 = 0\%$). The corrected figure is shown in this erratum (Fig. 1). This error has not modified the results and conclusions of the meta-analysis.

Author details

¹Department of Critical Care, Intensive Care Unit, Hospital de Clínicas (University Hospital), Faculty of Medicine, Universidad de la República (UdelaR), Italia Ave. 14th Floor. 11.600, Montevideo, Uruguay. ²Department of Anesthesiology and Surgery, Duke University School of Medicine, Duke Clinical Research Institute, 2400 Pratt Street, Office: NP 7060, Durham, NC 27705, USA.

Received: 31 January 2017 Accepted: 31 January 2017

Published online: 27 February 2017

Reference

1. Manzanares W, Lemieux M, Langlois PL, Wischmeyer PE. Probiotic and synbiotic therapy in critical illness. *Crit Care*. 2016;20:262.

* Correspondence: Paul.Wischmeyer@Duke.edu

²Department of Anesthesiology and Surgery, Duke University School of Medicine, Duke Clinical Research Institute, 2400 Pratt Street, Office: NP 7060, Durham, NC 27705, USA




Fig. 1 Effect of probiotic therapy on Mortality (n = 17). Original forest plot after reassessing the effect of probiotics on mortality