

RESEARCH LETTER

Open Access



Early identification of patients at high risk of group A streptococcus-associated necrotizing skin and soft tissue infections: a retrospective cohort study

Tomas Urbina¹, Camille Hua^{2,3}, Paul-Louis Woerther^{3,4,5}, Armand Mekontso Dessap^{1,3,6}, Olivier Chosidow^{2,3,5} and Nicolas de Prost^{1,3,6*} 

Keywords: Necrotizing soft tissue infection, Group A streptococcus, Toxin, Intravenous immunoglobulins, Clindamycin

Dear Editor,

Necrotizing soft tissue infections (NSTIs) are a heterogeneous group of devastating diseases involving a wide variety of microorganisms and affecting different body areas. The need for individualized treatment strategies has been recently put forward in a prospective cohort study of 402 patients in which group A streptococcus (GAS) infections were associated with more frequent septic shock [1]. Early identification of patients with GAS-related NSTIs could prompt initiation of targeted interventions, including clindamycin and intravenous immunoglobulins (IVIg). These drugs might be associated with beneficial antitoxic properties, but the level of evidence supporting them remains low (clindamycin) or highly controversial (IVIg) [2, 3]. The only randomized clinical trial evaluating the effect of IVIg specifically in patients with NSTI could not demonstrate a benefit on a composite outcome of death and quality-of-life evaluation at 6 months [4]. As previously commented [5], only 15% ($n = 13/87$) of included patients eventually had a microbiologically proven GAS NSTI. This was a major limitation and early identification of patients with a high probability of GAS-associated NSTIs would thus be crucial for further studies evaluating similar interventions.

A secondary analysis of a retrospective cohort including 224 patients admitted to our center for NSTI between

2006 and 2017 was conducted [6]. In accordance with the most recent guidelines, only patients with surgically confirmed NSTI were included (i.e., macroscopic appearance of tissues during operation as swollen, dull gray with a thin, brownish exudate with or without necrosis). Admission characteristics and microbiological documentation based on surgical samples, blood cultures, or subcutaneous puncture were recorded. We compared patients with a documented GAS infection to other patients regarding admission characteristics. A multivariable logistic regression model was used to identify admission characteristics associated with a subsequent GAS documentation.

Among 224 patients, 60 (27%) had a GAS infection, which was monomicrobial in 39 (17%) cases. Overall, 134 (59.8%) patients were admitted to the intensive care unit during their stay, of whom 113 during the first 24 h. Ninety-one (41%) patients presented with shock (i.e., required vasopressors), and 89 (40%) required mechanical ventilation. Sixty days after admission, 51 (23%) patients had died, including 10 (17%) with GAS, and 41 (25%) with non-GAS infections ($p = 0.255$, Mann-Whitney test). Admission characteristics associated with GAS infections by univariable analysis were non-steroidal anti-inflammatory drug treatment before admission and leukocytosis as a continuous variable. Those inversely associated with GAS infections were immunodeficiency, the nosocomial onset of infection, and an abdominoperineal location (Table 1). After multivariable analysis, only immunodeficiency (adjusted odds ratio (aOR) = 0.29 [0.10–0.74], $p = 0.015$) and an abdominoperineal location (aOR = 0.06 [0.00–0.30], $p = 0.007$) remained associated

* Correspondence: nicolas.de-prost@aphp.fr

¹Service de Réanimation Médicale, Hôpitaux Universitaires Henri Mondor, Assistance Publique – Hôpitaux de Paris (AP-HP), Créteil, France

³Université Paris-Est Créteil Val de Marne (UPEC), Créteil, France

Full list of author information is available at the end of the article



Table 1 Admission characteristics associated with group A streptococcal documentation

	Available data	Overall (n = 224)	GAS ^a (n = 60)	Others (n = 164)	p (univariate) ^b	Adjusted OR ^c	p (multivariate) ^c
Demographical data							
Age, years, median (IQR)	224	64.00 [53.00–74.25]	60.00 [50.00–72.00]	65.00 [55.50–75.00]	0.083		
Male gender, n (%)	224	127 (56.7)	31 (51.7)	96 (58.5)	0.443		
Comorbidities, n (%)							
Diabetes mellitus	224	83 (37.1)	18 (30.0)	65 (39.6)	0.244		
Immunodeficiency	224	58 (25.9)	6 (10.0)	52 (31.7)	0.002	0.29 [0.10–0.74]	0.015
HIV infection	224	2 (0.9)	0 (0.0)	2 (1.2)	0.954		
Cancer	224	21 (9.4)	0 (0.0)	21 (12.8)	0.008		
Corticosteroids	224	36 (16.1)	6 (10.0)	30 (18.3)	0.197		
Obliterating arteritis of the lower limbs	224	24 (10.7)	5 (8.3)	19 (11.6)	0.651		
Liver cirrhosis	224	9 (4.0)	0 (0.0)	9 (5.5)	0.142		
Chronic kidney disease	224	25 (11.2)	4 (6.7)	21 (12.8)	0.293		
Chronic alcohol consumption	224	27 (12.1)	5 (8.3)	22 (13.4)	0.422		
Obesity	224	57 (25.4)	13 (21.7)	44 (26.8)	0.54		
Prior to admission							
Time from first symptom, days, median (IQR)	224	5.00 [2.00–9.75]	5.00 [2.00–7.25]	5.00 [2.00–10.00]	0.599		
Antibiotic treatment, n (%)	221	137 (61.2)	30 (50.8)	107 (66.0)	0.057		
NSAID use, n (%)	222	46 (20.5)	19 (31.7)	27 (16.7)	0.024	–	0.122
Presentation upon admission							
Nosocomial infection, n (%)	222	45 (20.1)	4 (6.7)	41 (25.3)	0.004	–	0.197
Abdominoperineal location, n (%)	223	38 (17.0)	1 (1.7)	37 (22.7)	< 0.001	0.06 [0.00–0.30]	0.007
Shock, n (%)	220	91 (40.6)	21 (35.6)	70 (43.5)	0.369		
Creatininemia, μmol/L, median [IQR]	210	112.50 [69.00–171.25]	123.00 [71.25–187.25]	109.50 [67.25–167.75]	0.571		
Uremia, mmol/L, median [IQR]	207	9.80 [5.25–19.00]	10.25 [5.45–18.02]	9.80 [5.20–19.10]	0.966		
Plasma bicarbonate, mmol/L, median [IQR]	193	22.90 [19.70–26.80]	22.70 [20.10–26.00]	23.00 [19.50–27.05]	0.943		
Blood leucocytes 10 ³ /mm ³ , median [IQR]	219	14.40 [9.50–21.60]	17.20 [12.35–22.50]	13.60 [9.00–21.00]	0.016	–	0.067
Platelets 10 ³ /mm ³ , median [IQR]	189	217.00 [153.00–329.00]	223.50 [181.25–312.50]	207.00 [144.00–344.00]	0.45		
Hemoglobinemia, g/dL, median [IQR]	215	10.70 [9.45–12.15]	11.05 [10.15–12.50]	10.60 [9.35–12.10]	0.171		
Arterial lactate–mmol/L median [IQR]	146	2.00 [1.30–3.48]	2.10 [1.50–3.60]	2.00 [1.20–3.40]	0.677		

Analysis among 224 patients admitted for necrotizing soft tissue infection. ^aGroup A streptococcal infection. ^bp values for univariate comparison of documented group A streptococcal infection vs others; Chi-squared test or Fisher's exact test were used for categorical data according to sample size, Mann-Whitney's test was used for continuous variables due to non-parametrical distribution. ^cp values and adjusted ORs from a logistic regression model assessing the relationship between admission characteristics and group A streptococcal documentation. The model included all variables with a p value < 0.05 in univariate analysis. Analysis regarding 213 patients (11 patients excluded for missing data on one of the variables of the model. Immunodeficiency encompassed active cancer, chemotherapy within the last 3 months, previous HIV infection whatever the AIDS status, the CD4 lymphocytes counts or the viral load, any immunosuppressive drugs including chronic systemic steroid treatment (whatever the dose but for at least 3 months), HIV human immunodeficiency virus, NSAID non-steroidal anti-inflammatory drug

with the absence of GAS infection (Table 1). A sensitivity analysis using “monomicrobial GAS NSTI” as the dependent variable yielded similar results, except for younger age that remained in the model after adjustment (data not shown). Immunodeficiency ($n = 58$) and an abdominoperineal location ($n = 38$) had respective positive predictive values for the absence of a GAS infection (both mono- or polymicrobial) of 90% [79–96] and 97% [86–100] (Fig. 1).

In conclusion, we retrospectively identified two simple and available upon admission clinical predictors of GAS documentation among a large cohort of surgically

proven NSTIs. Our results show that NSTI patients with pre-existing immunodeficiency or an abdominal infection have a low probability of GAS infection and might thus not be suitable for inclusion in a trial assessing the effect of GAS-specific interventions. Such findings need to be assessed in a validation cohort in order to reinforce their generalizability. Improving identification upon admission of a subgroup of patients with a higher prevalence of GAS infection might help design future prospective trials aimed at assessing personalized treatment strategies [2].




Fig. 1 Diagnostic performances of abdominoperineal location and immunodeficiency for predicting absence of group A streptococcal documentation. The three top pie charts represent the proportions of group A streptococcal documentation, abdominoperineal infections and immunodeficiency in the whole 224-patient population of surgically confirmed necrotizing soft tissue infections. The two bottom pie charts represent the proportion of group A streptococcal documentation in the subgroup of patients with abdominoperineal infections (bottom left chart) or in immunocompromised patients (bottom right chart). Diagnostic performances of an abdominoperineal location of infection and of immunodeficiency for predicting the absence of group A streptococcal documentation were calculated using a contingency table approach. Immunodeficiency encompassed active cancer, chemotherapy within the last 3 months, previous HIV infection whatever the AIDS status, the CD4 lymphocytes counts or the viral load, any immunosuppressive drugs including chronic systemic steroid treatment (whatever the dose but for at least 3 months). PPV, positive predictive value; NPP, negative predictive value; Se, sensitivity; Sp, specificity

Abbreviations

GAS: Group A streptococcus; IVIG: Intravenous immunoglobulins; NSTI: Necrotizing soft tissue infection; OR: Odds ratio; PPV: Positive predictive value; NPP: Negative predictive value; Se: Sensitivity; Sp: Specificity

Acknowledgements

The members of the Henri Mondor Hospital Necrotizing Fasciitis Group are Romain BOSCH, Cécile CHAMPY, Olivier CHOSIDOW, Nicolas de PROST, Nicola DE ANGELIS, Jean-Winoc DECOUSSER, Camille GOMART, Jean-Michel GRACIES, Barbara HERSANT, Camille HUA, Raphaël LEPEULE, Alain LUCIANI, Lionel NAKAD, Alain RAHMOUNI, Emilie SBIDIAN, Françoise TOMBERLI, Tomas URBINA, and Paul-Louis WOERTHER.

Authors' contributions

All authors were involved in the study conception and design and conducted the study on behalf of the Henri Mondor Hospital Necrotizing Fasciitis Group. TU and NdP collected the data, performed statistical analyses, and wrote the original draft. All authors were involved in interpreting the data and reviewing the final manuscript. All authors read and approved the final manuscript.

Funding

This work did not receive any funding.

Availability of data and materials

The dataset used during the current study is available from the corresponding author upon reasonable request.

Ethics approval and consent to participate

The study was approved by the Comité de Protection des Personnes Ile-de-France V on March 8, 2018 (reference #16165). Patients received information during hospital stay that data abstracted from their medical charts could be used for research purposes.

Consent for publication

Not applicable.

Competing interests

PLW declares having received lecture fees and conference invitations from MSD.

All other authors declare no competing interest for this work.

Author details

¹Service de Réanimation Médicale, Hôpitaux Universitaires Henri Mondor, Assistance Publique – Hôpitaux de Paris (AP-HP), Créteil, France. ²Service de Dermatologie, Hôpitaux Universitaires Henri Mondor, Assistance Publique – Hôpitaux de Paris (AP-HP), Créteil, France. ³Université Paris-Est Créteil Val de Marne (UPEC), Créteil, France. ⁴Laboratoire de Bactériologie-Hygiène, Hôpitaux Universitaires Henri Mondor, Assistance Publique – Hôpitaux de Paris (AP-HP), Créteil, France. ⁵Équipe EA 7380 Dynamyc, Unité de Formation et de Recherche (UFR) de Médecine - site Créteil, Université Paris-Est Créteil Val-de-Marne, Créteil, France. ⁶Groupe de Recherche Clinique CARMAS, Université Paris Est-Créteil, Créteil, France.

Received: 27 November 2019 Accepted: 16 December 2019

Published online: 21 December 2019

References

1. INFECT study group, Madsen MB, Skrede S, Perner A, Arnell P, Nekludov M, et al. Patient's characteristics and outcomes in necrotising soft-tissue infections: results from a Scandinavian, multicentre, prospective cohort study. *Intensive Care Med.* 2019;45(9):1241–51.
2. Darenberg J, Ihendyane N, Sjölin J, Aufwerber E, Haidl S, Follin P, et al. Intravenous immunoglobulin G therapy in streptococcal toxic shock syndrome: a European randomized, double-blind, placebo-controlled trial. *Clin Infect Dis.* 2003;37(3):333–40.
3. Carapetis JR, Jacoby P, Carville K, Ang S-JJ, Curtis N, Andrews R. Effectiveness of clindamycin and intravenous immunoglobulin, and risk of disease in contacts, in invasive group A streptococcal infections. *Clin Infect Dis Off Publ Infect Dis Soc Am.* 2014;59(3):358–65.

4. Madsen MB, Hjortrup PB, Hansen MB, Lange T, Norrby-Teglund A, Hyldegaard O, et al. Immunoglobulin G for patients with necrotising soft tissue infection (INSTINCT): a randomised, blinded, placebo-controlled trial. *Intensive Care Med.* 2017;43(11):1585–93.
5. de Prost N, Lipman J, Mimoz O. Therapeutic targets in necrotizing soft tissue infections. *Intensive Care Med.* 2017;43(11):1717–9.
6. Urbina T, Hua C, Sbidian E, Bosc R, Tomberli F, Lepeule R, et al. Impact of a multidisciplinary care bundle for necrotizing skin and soft tissue infections: a retrospective cohort study. *Ann Intensive Care.* 2019;9(1):123.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

